

19

MANT

no waste
ht only a

stic. Hum-
t.


47
s are used
ed gallon
is used a
ere's low

and flow
waste. A
with each
d easy.

ESS

maximum
early sam-
ple ap-
mils, and
mish as
was done
that it is
then any
e.





THE GREATEST TRACTION TIRE EVER BUILT FOR FARM USE

THE Firestone Ground Grip Tire is without equal for traction. It is in a class by itself. In fact, the design and performance of this tire are so unusual that the United States Patent Office has granted a patent on it. Firestone provides one traction tread—Ground Grip—for cars, trucks, buses, tractors and farm implements.

Ground Grip Tires give your tractors greater drawbar horsepower, saving 25% in time and 25% in fuel. Your Firestone Implement Dealer or Tire Dealer has an economical plan for cutting down farm implement wheels, applying uniform diameter rims so a few sets of Firestone Tires fit all implements. Start today putting your farm on Firestone Ground Grip Tires and save time and money.

If you are ordering a new tractor have it delivered on Firestone Ground Grip Tires—the greatest traction tires ever built for farm use.

Listen to the Voice of Firestone featuring Richard Crooks—with Margaret Speaks, Monday evenings over Nationwide N. B. C. Red Network

Firestone

GROUND GRIP TIRES

FOR CARS • TRUCKS • TRACTORS • AND ALL WHEELED FARM IMPLEMENTS

Copyright 1937, Firestone Tire & Rubber Co.

APRIL,

High Color • No Worms • No Stings

Use **GRASSELLI**
ARSENATE of LEAD

The Mark of Protection

FOR MORE CLEAN, SOUND

High-Color **FRUIT**



THE recognized quality and effectiveness of GRASSELLI* Arsenate of Lead has made it the preference of critical fruit growers and orchardists in the heavily infested codling moth areas.

Its high strength, quick killing power, economy of operation, safety to fruit and foliage with high deposit, make it THE lead for the job.

When used with summer oils it acts both as an insecticide and

an ovicide offering additional protection against the codling moth.

GRASSELLI Arsenate of Lead goes into natural suspension in the spray rig, eliminating costly delays due to sludge, clogged screens and nozzles and blocked lines.

For Quality and Dependability choose GRASSELLI Arsenate of Lead this year.

Write for Grasselli Growers' Guide and Spray Chart.



OTHER GRASSELLI PRODUCTS—Bordeaux Mixture, Calcium Arsenate, Dutox,* Dry Lime Sulphur, Grasselli* Spreader, Lime Sulphur Solution, Loro,* NuRexform, Paradichlorobenzene, Sulphate of Nicotine, Sulfuron,* Zinc Sulphate and many others (*Trade Mark Reg.)

EXTRA

Orchard Brand News

EXTRA

VOL. 1, NO. 2

APRIL

1937

EASTERN EDITION

ARSENATE OF LEAD WITH "ASTRINGENT" GIVES BETTER CONTROL AT NO EXTRA COST

Quick Action Reduces Stings In Number and Size

In the face of increasing losses to the fruit fly, many commercial growers have turned to the Federal Government, State Agricultural Agencies, and various organizations for help in developing methods to reduce the damage caused by this pest.

With the use of the material known as "Astringent," the fruit fly is now under control and the number of stings is reduced to a minimum.

Since the fruit fly is a pest which attacks the fruit at an early stage, it is essential that the control measure be applied as early as possible.

percentage of full size worms were observed. The control of the fruit fly is a matter of timing, and the use of "Astringent" gives the best results.

Despite the fact that the fruit fly is a pest which attacks the fruit at an early stage, it is essential that the control measure be applied as early as possible.

Since the fruit fly is a pest which attacks the fruit at an early stage, it is essential that the control measure be applied as early as possible.



Apple Bill says, "Astringent" will provide more downright discouragement for worms than any other arsenate of lead."

Forms Close-Knit Uniform Film

Strong coverage on apples is obtained by lead arsenate in combination with "Astringent." The material is applied to the surface of the fruit, forming a close-knit uniform film.

After spray material is applied to the fruit, it is essential that the material be applied as early as possible.

Since the fruit fly is a pest which attacks the fruit at an early stage, it is essential that the control measure be applied as early as possible.

Since the fruit fly is a pest which attacks the fruit at an early stage, it is essential that the control measure be applied as early as possible.

Orchard Brand Arsenate of Lead with "Astringent" is each year consistently establishing new control records. Crop reports from growers throughout the country indicate an average of 15% to 20% increased codling moth control with "Astringent" over the ordinary arsenate of lead.

The high efficiency of "Astringent" lies not only in its rapid toxic action for killing more worms but also in the constant improvement of "Astringent's" physical properties which tend to make a tighter, firmer, more uniform cover on the fruit. Worm stings are drastically reduced, both in size and number, because "Astringent" protects the

complete surface and blocks the worm *before* it enters the fruit — not when the worm is half-way through the apple!

Because of its natural adhesiveness and "filming" quality, Arsenate of Lead with "Astringent" gives a greater degree of protection *with a minimum interference with leaf functioning*. Send today for information on

ORCHARD BRAND
REG. U. S. PAT. OFF.
"Astringent"
ARSENATE OF LEAD

GENERAL CHEMICAL COMPANY

Executive Offices: 40 Rector Street, New York City

Sales Offices: Atlanta • Baltimore • Boston • Buffalo • Charlotte • Chicago • Cleveland • Denver • Houston • Kansas City • Los Angeles • Minneapolis • Montezuma (Ga.) • Philadelphia • Pittsburgh • Providence • San Francisco • St. Louis • Wenatchee • Yakima • In Canada: The Nichols Chemical Company, Ltd. • Montreal • Toronto

PRODUCTS

LIME SULPHUR SOLUTION • DRY LIME SULPHUR • BORDEAUX MIXTURE (POWDER AND PASTE) • DRITOMIC SULPHUR • APPLE DRITOMIC SULPHUR • "ASTRINGENT" ARSENATE OF LEAD • STANDARD ARSENATE OF LEAD (POWDER AND PASTE) • BASIC ARSENATE OF LEAD • CALCIUM ARSENATE • ARSENITE OF ZINC • ZINTOX-A BASIC ZINC ARSENATE • PARIS GREEN • PARA-DICHLOROBENZENE • NICOTINE SULPHATE (40% NICOTINE) • X-13 (PYRETHRUM EXTRACT) • SULPHUR, ARSENICAL, NICOTINE, BORDEAUX, COPPER-LIME, AND ROTENONE DUSTS, IN DESIRED MIXTURES • INSTANT BLUESTONE • SOIL SULPHUR

GENERAL CHEMICAL COMPANY

40 Rector Street, New York City

Please send me information on Orchard Brand Arsenate of Lead with "Astringent," also a copy of the 1937 edition of "Cash Crops" Booklet. Also information on (see list of products):

Name _____

Address _____

City _____

State _____

AFG-2

Profitable PICKING DEPENDS UPON *Timely* SPRAYING



THERE'S a best time for every orchard operation—and you'll agree there's no more important a time than spraying time—a time when just a few hours may make for crop success or crop failure.

But the orchardist with a Cletrac can't be stopped, because Cletrac will go into the orchard any day or night, regardless of soil or weather conditions . . . up hill and down vale . . . through mud . . . around long turns or short turns . . . always with power enough to pull the spray rig under full load.

And when it comes to plowing, harrowing, discing or such odd jobs as stump pulling and hauling, Cletrac does them all efficiently because Cletrac is the all-year, all-job crawler

tractor designed especially for the fruit grower.

Here is a tractor that is fully streamlined, too, preventing damage to low hanging fruit and branches. You'll also appreciate Cletrac's exclusive controlled differential steering that keeps both tracks always pulling, providing positive control at all times between close-set trees and on sharp turns. Full vision forward helps the operator maintain a true course through any soil.

Study the many advantages of Cletrac that make it easier to crowd *more work—better work—timely* work into every working day and you'll wonder why you have so long been satisfied not to own a Cletrac.

THE CLEVELAND TRACTOR COMPANY • Cleveland, Ohio

Cletracs are available in sizes from 22–94 horsepower. Model E Cletracs use regular 70 octane gasoline or may be equipped to use kerosene. Larger models offer the choice of gasoline or Diesel power.

The Cleveland Tractor Co., 19310 Euclid Ave., Cleveland, Ohio

Send me your fully illustrated and detailed folder covering all uses of your streamlined orchard model.

Name _____

Address _____

Town _____

State _____

CLETRAC CRAWLER TRACTORS



Farm hands that pay their way

GOOD YEAR

IMPLEMENT TIRES

- Ask your Goodyear Dealer about the change-over plan which enables you to buy one set of Goodyear Implement Tires for use on several implements.



GOOD YEAR

TRUCK TIRES

- To get top mileage on truck tires choose the Goodyear type best suited to your particular hauling jobs. Your Goodyear Dealer can be of genuine service in helping you make your selection.



GOOD YEAR

AUTO TIRES

- The cheapest thing on your car is the best tire you can buy. Out of the experience of more than 21 years, car owners operating under all conditions have found there is no tire like a Goodyear Tire.



GOOD YEAR

KLINGTITE BELTING

- Goodyear Klingtite Endless Farm and Tractor Belting for heavy drives. Goodyear Klingtite Red and Pathfinder Belting, cut to length, for small power machinery. Goodyear Belting lasts for years, requires no dressing, works in any weather.



GOOD YEAR

BATTERIES

- There are four types of Goodyear Batteries in a variety of sizes covering a wide range of prices. In every one you get genuine Goodyear quality.



PAY FOR YOUR TRACTOR

with

GOOD YEAR TRACTOR tires

• SPECIFY GOODYEAR All-Traction Tractor Tires for your new tractor. Or cut down the steel wheels on the tractor you now have and fit them with Goodyear Tires.

The savings you make during the lifetime of the tractor will be greater than the tractor's original price.

Cutting down steel wheels to fit Goodyear Tractor Tires is a simple operation. Your Goodyear Dealer will tell you how it is done, where you can have it done and how little it will cost.

THE GOODYEAR TIRE & RUBBER COMPANY, INC., AKRON, OHIO

And once you make the change, you can count on savings such as these immediately: fuel, 25 per cent; labor, 33 $\frac{1}{3}$ per cent; oil and grease, 33 $\frac{1}{3}$ per cent; ordinary repairs, 40 to 50 per cent; depreciation 20 to 30 per cent.

In the meantime you enjoy the greater operating flexibility and the vastly increased comfort which pneumatic tires will give you.

Now is the time to put your tractor on rubber and start to save. See your Goodyear Dealer without delay.

GOOD YEAR

Money Savers on the Farm

AMERICAN FRUIT GROWER

TITLE REGISTERED IN

U.S. PATENT OFFICE

APRIL

1937

VOL. 57

THE NATIONAL FRUIT MAGAZINE

NO. 4

DIVERSIFICATION ON THE FRUIT FARM

IN the March issue of *American Fruit Grower* a series of articles by competent authorities on small fruit and their culture was offered to the readers, and in this issue a series is being presented on the stone fruits. The fruit industry of America is specialized and each group merits special consideration from time to time.

While statistics show that there are more individuals engaged in growing the apple than any of the other deciduous fruits, yet the others represent large investments and are often very hazardous because they are less hardy and, also, they have some serious insect and disease pests. This makes it all the more important to bring to our readers the latest information regarding them.

Diversification on the fruit farm is highly desirable in some sections and undesirable in others. So long as the erratic nature of our climate results in losses by frosts, freezes, hail, disease, and insects, the provident growers should have a cash income from more than one source or crop. But diversification of crops means diversification of information and techniques. This has been a stumbling block to many because it is not much of a human trait to change one's occupation or routine. But the important thing is to make the land earn all the income possible so far as it is consistent with good husbandry.

Probably one of the greatest stimuli to diversification is nearness to large consuming centers. The further removed one is from the large markets, the more likely he is to grow one or two crops on a large scale. The roadside and nearby city markets call for products over practically the entire year and this demand has resulted in many a grower changing his crop program to meet the demand.

Another advantage of diversification is the better distribution of labor. When the peak of the work for different crops comes at different times of the year, the labor can be employed more regularly, which is better for all parties concerned. We see this in many lines of agriculture. In the past the florist grew nothing but flowers and the vegetable forcer nothing but vegetables. Today his plant is often used to better advantage by growing some of both or all of one at one time of the year and all of the other at some other season. Similarly, there have been shifts from the large scale truck gardener to market gardening, and marketing to small fruit culture.

In the older communities a "balance" has been established between crops best adapted to the locality and conditions and the labor supply. Usually this should be departed from with caution. But there

are so many conspicuous examples of individuals who have changed their set-up and increased their income that it is worthy of careful study.

It is important to recognize the seasonal activities of the major crop and make any others fit into the off seasons. For instance, the critical labor period for apples begins in late August and pyramids until mid-October and then drops off perceptibly until mid-November. There is another smaller peak beginning about the first of April and extending during the spray season and spring work until early in June. On the other hand, there are comparative lulls during July and August, depending somewhat upon the amount of thinning, and again from December until mid-March. These seasonal periods can be recognized for each fruit and crop by those who keep books of their farming operations, or by the payroll accounts alone. This gives the best index of what other crops will fit into the orchard scheme with least interference.

True, many orchardists take advantage of these lull-times to get away from the farm and hold that that is an advantage over the live stock farmer. We have no quarrel with these individuals but not all prefer this option.

The best system of diversification would vary for different sections. That small fruits and stone fruits can frequently augment the income of the apple orchardist, and vice versa, must be patent to anyone who has seen the trend in many states. This is somewhat of a swing of the pendulum back to the earlier days, but a healthy one. With improved varieties of peaches, prunes, raspberries, strawberries, and others,

(Continued on page 36)

CONTENTS

Profitable Peach Production Factors.....	9
By F. P. Cullinan	
Peach Thinning or Tree Conditioning the Crop	11
By M. J. Dorsey	
"Little Leaf" Control of Cherries.....	12
Cold Hits Citrus Groves.....	12
Black Raspberry Mulching.....	12
Root Area of Strawberries.....	12
Certain Phases of Apricot Culture.....	13
By Warren P. Tufts	
My Forty Years in Plum Production.....	14
By Hiram Burkholder	
Into the Future with Eastern Cherries.....	15
By Carl G. Wooster	
Pollination with Particular Reference to the Grape—Part II.....	16
By Fred E. Gladwin	
Parafilm? Pliofilm? Phenothiazine?.....	17
By E. G. K. Meister	
American Pomological Society.....	18
A Page Conducted in the Interests of the Society	
State News from Near and Far.....	20
Cameral	22
Grape Gleanings.....	24
Control of Cherry and Plum Black Knot.....	28
By M. B. Cummings	
Nut Varieties.....	36
New Time and Money Savers.....	37
By Handy Andy	
Successful Orchards.....	38
A "Round Table" Page for Every Grower	
AMERICAN FRUIT GROWER	

AMERICAN FRUIT GROWER

Published Monthly by

AMERICAN FRUIT GROWER PUBLISHING CO.
1370 Ontario St., Cleveland, O.

SUBSCRIPTION RATES

Domestic, Except Cleveland, 3 years \$1.00. 1 yr. 50c.
Cleveland and foreign (except Canada) \$1.00 per year.
Canada, 50c per year.

Entered as second-class matter at Post Office at Cleveland, Ohio, under the Act of March 3, 1879. Additional entry at Mount Morris, Illinois.

ROGER FISON, EASTERN MGR.,
Room 1212, 30 Rockefeller Plaza
Phone—Circle 7-1863

NATIONAL ADVERTISING REPRESENTATIVES: J. C. Billingsley Co., 123 W. Madison St., Chicago, Phone Central 0465; 415 Lexington Ave., New York City, Phone Murray Hill 2-3912.

APRIL, 1937

E. G. K. MEISTER

Publisher

DEAN HALLIDAY

Managing Editor

EDNA ANNE KRAUSE

Associate Editor

T. J. TALBERT

Contributing Editor

DR. J. H. GOURLEY

Associate Editor

WILLIAM H. ZIPP

Field Editor



MARY LEE ADAMS

Home Economics Editor

PAGE 7

McCORMICK-DEERING TRACTORS BURN DISTILLATE

with UNEXCELLED EFFICIENCY



Here are two views of the popular McCormick-Deering Model O-12 Orchard Tractor. It is shown above equipped with special citrus fenders and at the left with standard equipment. The Model O-12 is a handy, compact, powerful little tractor that fits in well in most orchard, grove, and vineyard operations.

**But Fuel Economy Is Just One Thing
... You Also Get Performance, Long
Life, and Driver Comfort in These
Modern Tractors**



Above: Disking a young orchard with a McCormick-Deering outfit. TracTractors are available in three sizes: Model T-20 (shown), Model TA-40, and Model TD-40 Diesel.

McCormick-Deering Tractors earned the right, years ago, to be known as "balanced" tractors. In their design, no one feature has been stressed at the expense of other desirable features. For example, every McCormick-Deering Tractor is a true triple-power tractor, capable of perfect performance at the belt, on the drawbar, and through the power take-off.

Every McCormick-Deering Tractor has a 4 or 6-cylinder engine that develops an abundance of smooth-flowing power . . . power that gets the work done on time, and saves wear and tear on the tractor, machine, and driver.

McCormick-Deering "balanced design" goes all through the tractor, with every practical automotive development made use of to improve performance, economy, durability, driver comfort, and safety. This has made McCORMICK-DEERING by far the best buy in the tractor field, whether you need an all-purpose tractor, a regular wheel tractor, or an accessible crawler.

Phone the McCormick-Deering dealer for a demonstration . . . or ask us for full details.

INTERNATIONAL HARVESTER COMPANY
(INCORPORATED)

606 So. Michigan Ave.

Chicago, Illinois

McCORMICK-DEERING TRACTORS



Site selection is an important consideration in peach planting expansion. Trees in depression, shown in photo, are small and defoliated, while those on higher land show good growth.

PROFITABLE PEACH PRODUCTION FACTORS

By F. P. Cullinan

Senior Pomologist, U. S. Department of Agriculture

ACCORDING to figures recently released by the Bureau of Agricultural Economics, the number of peach trees in the United States has declined during the period of 1930 to 1935 from 79,000,000 to approximately 67,000,000, and bearing trees from 59,000,000 to 54,000,000. The proportion of trees not of bearing age has declined from 25 per cent to 19 per cent of the total. It is expected that the annual production of peaches during the next five years will be somewhere around 45,000,000 to 50,000,000 bushels. In other words, unless growing conditions are extremely favorable, with the present number of bearing peach trees we should not expect the peak of production of 76,500,000 bushels reached in 1931. This was the largest total production in this country in 36 years.

With this decrease in the number of bearing trees and the prospects for

The following article is a digest of Dr. Cullinan's talk presented before the joint meeting of the American Pomological Society and the Virginia State Horticultural Society.—EDITOR.

continued fair prices for peaches during the next five years, it is probable that there will be considerable new replanting. It would be most unfortunate, however, if there should be a marked increase above what is necessary to replace trees now going out of production.

Factors that should be considered


in a rational replanting program include site in relation to frost hazard, varieties that have proved unprofitable because of unproductiveness or susceptibility to cold injury, plantings in regions where severe winter injury has resulted in heavy bud and tree mortality, varieties that have proved unpopular on the market, and the production of poor quality peaches that are not only unprofitable to the producer but tend to lower market prices.

Although the prices of peaches during the past two years have been very satisfactory for those who have been successful in obtaining a good crop, the profitableness of the business

"It is a safe prediction that commercial peach varieties of the future must be of higher quality if this fruit is to hold its place on the market in competition with other fruits"—

Says Dr. F. P. Cullinan, who is shown here as the American Fruit Grower camera caught him just after delivering his address, reviewed on this page, before the fruit meeting of the American Pomological Society and Virginia State Horticultural Society at Roanoke, Va.





Experiments are being carried on to reduce the amount of cultivation necessary in the peach orchard as is being done in above photo at the Mantle and Mantle Lake Erie Farms Orchard in Ohio.

should be considered over a longer period, based upon the average yield per acre obtained during the years of crop failure as well as the years of good production. Most fruit growers, both apple and peach, are convinced after the past six or seven years' experience that they cannot hope to remain in the fruit business if a high average yield per acre is not maintained.

The problem of maintaining vigor in peach trees is one of the most important to consider. The peach tree, relatively speaking, is not a long-lived tree, nor is it especially vigorous and hardy except under the most favorable conditions. The response from applications of nitrogen is particularly noticeable when the trees are young and have not produced heavy crops of fruit. As the trees grow older it becomes more difficult to maintain the proper terminal growth and vigor in peach trees in years following heavy production. On some soils we find that peach trees do not continue to maintain satisfactory terminal growth even with fairly large applications of nitrogen. It is not to be inferred that

nitrogen is the only fertilizer element that is beneficial to peach trees. We know that trees in order to grow and be productive require other fertilizer elements, such as phosphorus, potash, magnesium, etc. These, however, have usually not been of great concern to the fruit grower in the past since the soils themselves, generally speaking, have been fairly well supplied with adequate reserves of these mineral elements or they could be added through cover crops or barnyard manure.

The use of cover crops as a means of increasing the reserves of nutrient supply in the soil has not been as widely practiced by the peach grower as it deserves to be. During the past four years we have been carrying on some studies to determine whether or not we can satisfactorily grow a young peach orchard with a minimum amount of cultivation. We have had in mind two things: first, the increase in the fertility of the soil through the use of cover crops, with the possible increase in organic matter content, and, second, the effects of these cover crops on moisture and nitrates as they

affect growth and production. The trees received uniform cultivation during the first year, and a summer cover crop of soybeans was grown followed by rye and vetch in the winter. Beginning with the second year, we have used on the different plots sweet clover and lespedeza as legume sods, summer covers of soybeans, buckwheat and crotalaria, followed by a winter cover crop of rye and vetch.

The lespedeza plot was disked up thoroughly in late winter to incorporate the dried plants and seeds into the soil, and thus facilitate reseeding. A fine stand of plants was obtained every year by late March to early April. The sweet clover was seeded in March, when plots were first established, and was not clipped in this or the following year. With the second biennial crop, it was clipped in the first and second years, being disked up after seed formation in the second year, and seeded again in August. In the four years of this experiment there have been two crops of biennial sweet clover.

Where summer cover crops of soybeans and buckwheat were used they were seeded about June 1. The crotalaria was usually seeded two or three weeks later. The summer covers were disked up in late August and were followed with a seeding of rye and vetch for a winter cover. This in turn was disked up in mid to late April, depending upon the earliness of the season.

At the time of seeding of the summer and winter cover crops, 250 pounds per acre of a 5-8-5 fertilizer was drilled in with the seed. A similar amount was drilled in the lespedeza and sweet clover plots early in the spring. This fertilizer was primarily for the growth of the cover crops and the trees received what was considered an adequate application of nitrogen consisting of one-half pound of nitrate of soda in the first year and increased to three pounds in the fifth year, broadcast around the tree.

The best growth to date on the five-year-old trees has been made on the plot receiving only a winter cover of rye and vetch. The trees receiving summer covers, particularly soybeans, made very satisfactory growth but it was apparent that they were somewhat smaller. The trees in sweet clover and lespedeza were the smallest of all.

In the four years of this experiment, particularly in 1934, 1935, and 1936, we had periods of dry weather during the growing season and were thus afforded an opportunity to study the effects of the cover crops on moisture.

(Continued on page 34)

Organic matter is of importance in peach soils. Illustration on the left shows Joe B. Hale, secretary of the Illinois State Horticultural Society, preparing to spread straw in his peach orchard at Salem, Ill.



PEACH THINNING

OR TREE CONDITIONING THE CROP

By M. J. Dorsey
University of Illinois

THINNING is a very necessary part of peach growing and takes its place primarily as a means of regulating the size of fruit. Fortunately, thinning is not always necessary, but when it is, it is almost fatal to leave this operation undone. Since the more recent research shows that thinning is not as simple as it was once thought to be, let us take a practical look at the thinning problem and take stock of some of the variables involved in tree conditioning the crop.

The peach grower is confronted with a thinning job any time the bud survival is sufficiently heavy for an excess set. Potentially, this is always the case, because a mature peach tree may bear anywhere from 5,000 to 40,000 flowers. Counts have frequently been made of trees bearing as many as 4,000 to 6,000 fruits after the June drop when a further reduction in the number is relatively light. Consequently as the season advances, the problem of weight alone becomes more and more serious.

The question, then, of how many fruits should be left on a tree becomes a very important matter. As an approach to the problem, we might well consider, first, the relationship between size of fruit and the number per bushel. Counts in the Illinois experiments show that the number of fruits of the different sizes in 50 pounds to be as follows: 1¾ inches down, 500; 1¾ to 2 inches, 300; 2 to 2¼ inches, 240; 2¼ to 2½ inches, 210; 2½ to 2¾ inches, 150; 2¾ to 3 inches, 100.

With these data at hand, it will be seen at once that there are two ways of obtaining yield per tree: one by increasing the number of fruits and the other by increasing size. Naturally, then, the thinning problem takes its place in "adjusting the crop to the tree," to use an old expression, somewhere near the point where the better commercial sizes will be reached under average conditions.

Experienced growers can tell at a glance what would be a fair yield for a given tree condition year in and year out. For our present purpose, suppose we assume that five bushels would be a fair yield for a given tree condition. Obviously, from the number per size given, five bushels may contain as many as 2,500 peaches or

as few as 500, depending upon the size. If the desired size for a given tree condition would be about two and one-fourth inches, then it will be seen that in the neighborhood of 1,000 or 1,200 peaches should be left on the tree after thinning.

Of course, it is out of the question to count all of the peaches in an orchard, but in starting crews to work it is easy to pick out a typical tree, count the fruit on it before thinning, thin it, and then count the number pulled off as well as the number left on. Experienced help will soon get the pattern in mind and the crop can be evened up in the orchard very skillfully, leaving everything on some limbs or thinning others according to the type of tree. In this way thinning can be done without seriously cutting the yield, although our experience shows that it is difficult to determine exactly the number of peaches which should be left on the tree to get the maximum size for a given tree condition without reducing the yield.

With the potential set so heavy and the thinning job increased accordingly, the time of thinning then becomes a very important consideration. Under practical conditions the practice heretofore has been to wait until after the June drop. This recommendation needed experimental verification however, so at the Illinois station the results show that the time of thinning can be left without serious risk until after the June drop.

This whole matter can be readily understood by taking into consideration the natural drops on one hand and the growth periods of the fruit on the other. The usual drops in fruit are referred to as the first, second, and third or June drop. This latter is also sometimes spoken of as the physiological drop. It is by far the most conspicuous of the three in the peach, because the fruit is larger at the time, although the reduction in the total number of blossoms or fruits may be extensive in either of the first two drops. It would seem unnecessary to start thinning before the natural drops are over unless the acreage would require it.

The measurements at a number of stations have demonstrated three well defined growth periods in the peach.

(Continued on page 29)

AMERICAN FRUIT GROWER



About 100 fruits in this bushel.



About 240 in this one.



And 400 in this one. How much difference in price?



A good job of thinning as shown by the uniformity of these "tree run" peaches.

PAGE 11

CHERRIES •

"LITTLE LEAF" CONTROL

Sweet cherries suffering from the condition known as "little leaf" have dwarfed, light-colored foliage which is often clustered. A tree affected in this manner usually dies in a year or two or is so weakened that it is worthless. This condition is also known as "rosette" and may be troublesome on many deciduous fruits.

While speaking before the Oregon State Horticultural Society, O. T. McWhorter, Oregon extension horticulturist, recommended the use of zinc for the improvement of trees affected with "little leaf." He stated that zinc tacks were suggested for small trees. The tacks are placed one-half inch apart and spirally around the tree. They are spaced one inch or more apart around the trunk to prevent girdling. Placing of the tacks is important, and they should be driven into the tree at least one-half of their length.

On badly affected trees of all ages at The Dalles, zinc sulphate used at the rate of one-half pound to 50 gallons of water, with or without lime, gave satisfactory results. When lime is desired, it is used at the rate of one-twentieth pound to one pound of zinc sulphate. Less injury to foliage was noted at The Dalles when spraying was done after 4:00 in the afternoon. The "little leaf" condition on young cherry trees, causing dwarfing and usually death of the tree, merits attention by careful use of zinc sprays and zinc tacks until the trees reach seven or eight years of age and are free of all symptoms of the condition.

Mr. McWhorter indicated that from available information the first zinc spray should be given just before the buds open and others should follow a month after the leaves appear. Zinc sulphate sprays using 10 to 12 pounds to 100 gallons of water with five pounds of lime have given good results where there is danger of foliage damage.

Zinc sulphate injections were made in trees eight years of age and older by boring holes an inch and a half deep, three-eighths inch in diameter and spaced four to five inches apart around the tree trunk at the ground line. The holes were filled with zinc sulphate to within one-half inch of the outside. The openings were then sealed with grafting wax.

Cherry trees which had 90 per cent affected leaves have shown normal appearing foliage of a dark green color the following year after zinc injections were given in the fall. July-treated trees showed improvement in six weeks with no sign of injury resulting from the injections.



A heavily mulched raspberry planting

CITRUS •

COLD HITS GROVES

California citrus growers are contemplating tree repair measures after a period of extreme temperatures never before equalled in southern California. While much fruit was frozen during the cold wave, it was surprising to note that there was comparatively little tree splitting.

Orchard heating practices stood a most grueling test against the low temperatures. It is estimated that 80,000,000 gallons of oil were consumed in the heaters. Cost of the oil alone has been set at \$3,200,000 and it is probable that total cost of heating practices for grove protection amounted to more than \$10,000,000. Growers who heated their groves were repaid by the survival of a good portion of their crops, while those who did not resort to heating lost their entire crop. It is stated that 4,000,000 heaters were used to combat this freeze.

Growers derived some benefit from the low temperatures in that future precautions will be taken to provide sufficient storage capacity for fuel oil, a greater supply will be carried throughout the danger seasons and facilities for rapid delivery of oil will be arranged. In the future growers will probably hire two crews of workers to take care of lighting and refilling the heaters and hauling oil.

California horticulturists advise that no citrus pruning be done during the coming season and that pruning next year should be light, confined to dead-wood removal, with thinning out where foliage is particularly heavy. For protection against sunburn, it is desirable to whitewash exposed branches. Unless disinfectants are applied to injuries before they are covered with a wound dressing, there is danger of heart-rot fungi attack. A good disinfectant is one quart denatured alcohol, one-fourth ounce mercuric cyanide and three quarts of water.

BERRIES •

BLACK RASPBERRY MULCHING

Results of a two-year test on black raspberry mulching, conducted by Dr. Leon Havis of the Ohio Agricultural Experiment Station, have recently been announced. This test was started in March, 1935, when wheat straw was applied at the rate of five to six tons to the acre. The plants were Cumberland black raspberries set out in 1928. Unmulched check plots were cultivated during the two seasons of the test.

Value of the straw mulch was especially significant in 1936. Foliage, vigor of new canes, and general appearance were superior in the mulched plots. In 1935, the total number of canes produced was only slightly higher in the mulched plots, but in 1936 the number produced was much greater.

Other advantages of the mulching were apparent. Berries grown in mulch are less likely to be hampered in appearance by soil contact. There is no need for cultivation of the mulched plot as weeds are almost entirely eliminated. The prevention of soil moisture loss is another definite advantage of mulching. In addition to these benefits, Dr. Havis states that evidence indicates the mulch aids in protecting the roots and crowns from winter injury.

Labor involved in applying the mulch may be spread over a long period of time and can be done when the grower is least busy. Danger of fire from the mulch may be minimized by leaving strips of unmulched soil in the planting. In some cases only alternate strips between the rows are mulched when fire hazards are present.

It is evident that when wheat or oats straw is used for mulch, it is necessary to increase the usual application of nitrogenous fertilizers. This practice is especially desirable for the first two or three years after applying the mulch.

Where mulching is practiced only in the rows and a few feet on each side of them, it is necessary to cultivate the areas not mulched.

STRAWBERRY ROOT AREA

On some soils it has been found that 90 per cent of the total strawberry root system is located in the top six inches of soil. In many cases 70 per cent of the root system was in the first three inches. This feeding area indicates need for proper moisture in strawberry soils, supplied by irrigation, mulching, and the addition of humus materials.

Top (Fig. 1)—Eighty-year-old Royal apricot tree on peach root, pruned by the so-called Winters system. Pruning, thinning and harvesting are accomplished from a nine-foot ladder. This tree has a spread of more than 60 feet.

Center—Fifty-year-old Royal apricot tree on Myrobalan root. The system of pruning here employed is typical of possibly 50 per cent of the bearing apricot acreage of California. Note that bloom is largely confined to upper portions.

Bottom—Twelve-year-old Royal apricot orchard on apricot root, pruned by "thinning out" only. Trees are planted 25 by 25 feet. Orchard operations are conducted with a 10-foot ladder.



CERTAIN PHASES OF APRICOT CULTURE

By Warren P. Tufts

Pomologist, University of California

IN addition to the United States, only Australia, the Union of South Africa, Syria, and Persia produce apricots in commercial quantities. Comparable statistics are unavailable. Australia and South Africa sending canned and dried apricots to the markets of the United Kingdom, compete directly with the United States. In general, Syria and Persia do not enter competitive markets.


Commercial production in the United States is practically confined to California, which in 1930 had 93 per cent of the total tonnage and 89 per cent of the trees. Washington, ranking next, produced three per cent. Between 1920 and 1930, Utah, Colorado, Oregon, Idaho and Washington increased their plantings 560 per cent; California 13 per cent. In 1929, the 11 western states produced 98.6 per cent of the U. S. tonnage. The western states, other than California, market their apricots in the fresh condition; largely in the Pacific Northwest, which does not seriously compete with the California crop, which is sold dried (70 per cent), canned (20 per cent) and fresh (10 per cent). California has the earliest season (May 15 to July 15).

Before the World War, 68 per cent of the dried product was exported; during 1922-1926 only 53 per cent. More apricots are now canned than formerly and fewer dried. Although fresh shipments in California have increased in recent years, they constitute a relatively small portion of the annual crop of about 240,000 tons from 80,000 acres.

Most growers of deciduous fruits

(Continued on page 26)





The Stanley plum, one of Mr. Burkholder's favorites which he will use in his new planting.

MY FORTY YEARS IN PLUM PRODUCTION

By HIRAM BURKHOLDER

IF you are going to plant a plum orchard, I believe you will be interested in experiences I have had with plums over a 40-year period. When I planted my first trees in northern Ohio, I knew scarcely anything about kinds and so set the varieties widely offered at the time. Most of these proved unsatisfactory, the fruit being insipid or too acid. I have since dug out many of these trees and have grafted over many others.

Such promiscuous planting in nearly all orchards has been responsible for the comparatively light consumption of plums. Consumers have turned to other fruits after a few experiences with poor quality sorts. It is going to take years of growing and selling the better varieties to retrieve the plum market. Many opportunities for developing a worthwhile market and making a profit await the grower starting out at this time with quality varieties, in my opinion.

In the matter of varieties, let us first consider the Japanese sorts. These are early bloomers and are frequently killed by late frosts. They are clings and are acid about the pit

or skin or both. The foliage is tender and is injured by the sprays used on the European varieties. This is an important item where they are not planted in large blocks so that sprays can be mixed especially for them. The fruit of such types as Monarch, Gneii, Shipper's Pride, Moore's Arctic, Imperial Gage, Quackenboss and others is very apt to discourage consumers in the use of plums.

The old Lombard continues to be a favorite with some of my customers. It is good when thinned and allowed to ripen on the tree, but it is biennial and loads so heavily in the bearing year that it requires thinning to obtain size and quality. Imperial Gage is fine for eating fresh, but when cooked is soft and more acid than Reine Claude.

If stocks were wanted upon which to "work" better varieties of weak-tree tendencies, we could not do better than to use Imperial Gage, Hand and Quackenboss.

Of the varieties I am setting to increase our plantings, I would mention Stanley prune first. It is a good grower, an early bearer, of high qual-

ity, free stone, an annual producer, and ripens about with Italian prune.

Imperial Epineuse is a high quality prune, of reddish color, and one of the finest in the prune class to eat out of hand. It sells freely from roadside stands. It is also excellent when cooked. The tree has a scrubby growth for several years but later makes a good spreading tree. Its buds are hardier than Italian or Hall, but not quite as hardy as Stanley. It is not nearly as early in starting to bear as Stanley. Demands for this prune among my customers are increasing.

I must mention my experience with the new Hall prune. It has proved very susceptible to winter injury of the fruit buds. The tree forms nicely but is not a fast grower and is quite subject to shot hole fungus and body trouble near the ground, as is Grand Duke. It is one of the finest plums in shape and quality I have seen, but I am starting to work my youngest trees over because of the tenderness of the fruit buds.

Another of the new plums mentioned favorably in some places is the Albion. I have a few young trees not bearing but expect to plant more, although some growers report it as not being very prolific. The fruit is large and late ripening.

I have grown just a few Elephant Hearts on grafts. I am not familiar with the tree characteristics, but the plum is of good size and quality.

German plums make large, lovely shade trees but have not produced enough fruit in the past to be worth considering. The fruit is small but good. Italian prune (Fellenburg), of which I have more trees than any other variety, is late starting to fruit and seldom has a good crop.

I have a good many large French damson trees (my favorite damson). Shropshire damson fruit is smaller, of not as good quality, and the tree is not shaped as well. It sells as well as the French on city markets, however, because some consumers are suspicious that French is not a damson. I sell the majority of my damsons to hucksters bound for Columbus or farther south, and I also do some trucking to the south. Mining sections are a good outlet.

Reine Claude is my best seller in the orchard, outside of the prune class. It is not a good "city" plum as there are several other green gage types of inferior quality on the market and the average city consumer does not know which is the good one.

In a shipment of nursery stock which I purchased in 1918 were some Hudson plum trees. The fruit of these is very good and sells well to both hucksters and consumers.

Grand Duke, though not sweet, is one of my best sellers. Being large and attractive, hucksters and commission houses buy them readily. The tree is subject to body trouble,

(Continued on page 25)

INTO THE FUTURE WITH EASTERN CHERRIES

By CARL G. WOOSTER

OUR experience as cherry growers is very closely associated with the development of the industry in Wayne County, New York. About 1900, my father purchased a farm of 176 acres near Lake Ontario, which had on it at that time about 80 acres of apples and one acre of Napoleon sweet cherries.

For many years the returns from the one acre of sweet cherries could be depended upon to pay all the taxes on the farm. The cherries in those days were sold on the fresh fruit market. During the next few years the canning industry developed and with it came the demand for red sour

nearby markets for fresh fruit, nevertheless a rapid readjustment of plantings into a narrower range is taking place.

New York State is the second largest producer of red sour cherries. Wayne County produces about half of all cherries processed in the State, and the greater part of these are grown within two or three miles of Lake Ontario. An analysis of the varieties which our better growers are planting and the cultural methods employed would seem to answer many questions relative to the future of the industry.

About 70 per cent of all plantings

Windsor is somewhat later than either Napoleon or Yellow Spanish, but must be harvested before reaching full maturity if it is to be bleached for Maraschino purposes. Windsor is a hardy tree and bears abundantly when cross pollinated. It is also a very good cherry for the fresh fruit trade.

Schmidt should be mentioned in reference to fresh fruit. It is not surpassed in this respect, and does not ordinarily crack. It is somewhat slower in reaching bearing age, and is not acceptable to the Maraschino trade.

In selecting a site for the cherry



The author inspecting a tree in a 25-year-old cultivated New York Montmorency orchard planted on the 20 by 20-foot system. To the right of Mr. Wooster is Elmer G. Butts.

cherries. Later the freezing process was developed, and, more recently, the famous Maraschino process for sweets and also sour.

Prices were high for cherries and the demand exceeded the supply. Trees were set by the thousands, many of them without regard to soil requirements, nor to the limiting factor of frost protection which is afforded by our Great Lakes to a restricted area.

During seasons of favorable weather, many of these marginal orchards produce a crop and share in the low price paid for cherries. In less favorable years they do not bear. While the price level of cherries has dropped in the last few years, a more important factor in determining the profit in cherry growing is the ability of an orchard to produce a heavy annual crop. Thus, while some cherries of several varieties will continue to be grown in many parts of the country to meet the demands of local and

today are Montmorency. The tree is hardy, productive, and the fruit meets the demands of the consumer. Early Richmond is only second-class. English Morello is not in demand for processing or freezing. The trees are more fastidious as to soils, require heavy feeding, and the foliage is very susceptible to leaf spot.

There is some increase in the plantings of sweets due to the growing demand for Maraschino cherries. Napoleon, sometimes called "Royal Ann," is most popular at present. Windsor and a few Yellow Spanish are being set as pollinators. The Napoleon will crack if rains occur at harvest time. Yellow Spanish is not as susceptible to cracking, and can usually be harvested after Napoleon.

orchard, the naturally well-drained deep soils of porous nature are best. Cherry trees will not tolerate wet soils, and while tile drainage does help somewhat, nothing equals good natural drainage. Good air drainage should also be given consideration.

Montmorency, as well as the sweets, should be budded on Mazzard rootstock. The trees grow larger, are thriftier, bear heavier, and live much longer than on Mahaleb. We sometimes plant sour on Mahaleb when they are to be used as fillers, and for this purpose they do very well.

Montmorency on Mazzard roots, planted in good soil, should be set not less than 20 by 20 feet, and sweets not less than 30 by 30 feet. Some of our best Montmorency orchards are 24 by 24 feet on the square. These distances facilitate spraying, cultural, and harvesting operations, without bruising and destruction of

(Continued on page 30)



A close-up of the Fredonia grape, taken during 1936 season at New York Vineyard Laboratory. This superior variety was originated by F. E. Gladwin.

POLLINATION

WITH PARTICULAR REFERENCE TO THE GRAPE

By FRED E. GLADWIN

Viticulturist, New York Agricultural
Experiment Station

PART II

The cross-pollination and subsequent fertilization of grape blossoms is almost entirely subject to air currents, as they alone disperse the pollen. Bees or other insects play but a small part in the dissemination of grape pollen. Investigations show that grape pollen has been carried approximately 16 feet by gentle to fresh air currents, while sticky screens placed 24 feet distant from the pollen-bearing vine used in the tests failed to show any trace of pollen adhering to them.

These tests would indicate that if other varieties are to be interplanted for the purpose of cross-fertilizing grapes, the pollen-bearing vines should not be over 16 feet from the variety to be served. Further, the pollinators should literally surround the self-sterile and imperfectly-fertile varieties so that the full advantage of changing winds may be had.

Repeated tests have shown that many varieties may be made to bear more compact clusters through the artificial application of viable and compatible pollen. Concord and Niagara, which normally set satisfactory clusters, have been made to yield more berries per cluster when hand pollinated with pollens from other varieties. Butler and Lindley, two imperfectly-fertile varieties, were made to produce compact to very compact clusters when viable pollens were applied by hand.

In early work, the pollen was applied by two methods: either fresh blossom clusters of the pollen-bearing variety were dusted over the clusters of the varieties mentioned, or the blossom clusters were taken from the pollen-bearing vine, dried, finely pulverized by rubbing through the hands, and when a sufficient fining had been obtained, the powder, which included much pollen, applied to the blossoms of the varieties to be cross-fertilized as soon as the stigmas of the latter were seen to be receptive. For purposes of application, fine camel's hair brushes are very satisfactory. This method of cross-pollination is entirely practical on a small scale as in the home garden, but its application in a commercial way is not justified because of its time-consuming labor.

The following varieties (now quite generally grown) are strongly self-fertile: Clinton, Concord, Catawba, Delaware, Agawam, Diamond, Niagara and Worden. Some of the self-sterile and imperfectly self-fertile are: Barry, Brighton, Herbert, Lindley, Eumelan, Salem, Wilder and Vergennes. It has been stated before that most self-fertile varieties are greatly improved through cross-pollination, but for the self-sterile varieties cross-pollination is absolutely necessary to insure the production of marketable clusters.

Pollen that matures before the

stigmas are receptive may be dried, and if kept dry, held in good viable condition for several days, and sometimes a few weeks. Thus if both essential elements, ovules and pollen, do not reach the fusing stage coincidentally, pollen that matures before the stigma is receptive can be kept until such time. Since many varieties produce very viable pollen, there should be no difficulty in having an ample supply available throughout the entire range of blooming, from those that bloom very early and those that come to blossom very late.

Throughout all of the work in cross-pollination there has been one outstanding pollen-producing variety which has furnished pollen for most of the artificial applications. This variety is a male vine, hence sets no fruit. It was planted because of its possibilities as a graft stock. It annually produces a mass of very large blossom clusters, and it is one of the few that is decidedly fragrant. However, this pleasing odor attracts no bees and hence its pollen which is produced in great abundance is entirely disseminated by air currents.

For several years this variety has been used in the breeding of new grapes, but all of the seedlings have been males. All of the progeny have thus been indelibly stamped with the characters of the male parent. These trials indicated that here was a most viable pollen, and such it has proven to be whenever it has been used to improve the setting of varieties that ordinarily set straggly clusters.

The variety is listed as "Aramon crossed with Rupestris Ganzin No. 1." It can be found in most collections of grapes wherever any work with grape grafting has been done. It propagates readily from cuttings and grows quite vigorously. A few vines will furnish pollen sufficient to cross-pollinate several hundred vines. When it is planted not over 16 feet in any direction from a variety that is self-sterile, good results may be obtained without hand application.

During the past year pollen from "Aramon X Rupestris Ganzin No. 1" has been applied to the blossoms of a self-sterile variety, Eumelan, by means of a small hand duster such as is ordinarily used in dusting plants for insects and diseases. Eumelan is a variety highly prized in the manufacture of wine, but because of the poor setting of its blossoms it is but little grown.

As fast as the blossoms of this male variety opened, the clusters were picked from the vines, placed on screens in a room free from air currents and thoroughly dried. They were then pulverized, placed in tins and sealed. Two weeks later this pol-

(Continued on page 35)

PARAFILM

?

PLIOFILM

?

PHENOTHIAZINE

?

By E. G. K. MEISTER



YOU'LL FIND ALL THE ANSWERS IN THE COMING ANNUAL JUNE DIRECTORY EDITION AMERICAN FRUIT GROWER

NEW names and new products are constantly appearing in the business of horticulture. These new names are confusing to fruit growers and, being technical, are difficult to remember. The successful grower, however, is usually the first to apply new ideas and materials. He keeps an accurate record of new things that come to his attention and seldom depends upon his memory when a problem arises.

A part of the library of every successful grower is the annual Directory and Buyer's Guide Edition of AMERICAN FRUIT GROWER. It contains a complete index of anything and everything needed in orchard work. Many new products will appear in the revised edition, which is now being compiled and will be published in June. Such problems as the following need no longer puzzle fruit growers:

★

"Sorry, boss, but we can't go ahead with the spraying until we get a new nozzle for the gun. The old one's given out. Who makes the XX model?"

★

"Now that I've decided to build a cold storage after hearing so much on the subject at the last state horticultural meeting, wonder where that blower type refrigeration unit can be obtained."

★

"Let's see, what was the name of the company handling the new type pruning shears I saw demonstrated? Wish now I'd written down the name and address, so I could send for a pair."

How many times have you started your day with questions like those above puzzling and perplexing you? And haven't you wished every time that you had a handy buyer's guide to give you just the information you needed?

Another feature of the June Directory Edition, invaluable to all growers, is a complete and up-to-the-minute treatise on cold storage for the fruit farm. Cold storage of fruit was first attempted by Seneca, when ancient Romans collected snow from the mountains. Alexander the Great likewise dug trenches which were filled with snow and covered with straw and twigs to preserve the perishable fruits of that time.

Refrigeration by artificial means began in 1825 with the development

of an air pump to extract air from water. Early expensive machinery has been improved, simplified, and standardized, reaching its highest development in the small household electric refrigerator. Mechanical refrigeration provides growers with cold storages which are a sure way to high fruit prices.

"How to Convert a Common into a Modern Cold Storage" will be discussed in detail in the June Directory number. Another article deals with how to build a 5,000-bushel cold storage. Experienced authors will give specifications, construction details, and plans for building and equipping such a structure.

The story of modern mechanical refrigeration for fruit growers holds much of interest and value to growers.

WATCH FOR THE JUNE ISSUE CONTAINING COMPLETE COLD STORAGE DATA



A PAGE CONDUCTED IN THE
INTERESTS OF THE AMERICAN
POMOLOGICAL SOCIETY

PLANT WITH CARE—AND ADVERTISE

DECIDUOUS fruit growers in all sections of the United States and Canada are facing a new season with considerably more confidence than was manifested a few years back. The deciduous fruit industry as a whole marketed its crop for more nearly satisfactory prices than has been the case for some time. Apple prices have ruled firm and upward most all the season so that those regions and growers in particular who were fortunate enough to have a crop have felt the influence of a better price.

There is a feeling of optimism among apple growers. The number of bearing and non-bearing trees has been greatly reduced, and the surpluses of former years are not likely to menace the security of the industry for a number of years. There is a feeling that apple growers are now due for a series of years of relatively good prices.

Such a feeling, however, may again promote overplanting of the kind witnessed 20 years ago. Planting at this time is desirable and necessary to stabilize the industry, but careful observers feel the amount of plantings should be only large enough to make the acreage replacements which are needed to keep the supply abreast of the prospective demand.

The various institutes and the advertising campaigns which have been organized in various regions have in a few short years made a real impression on the consuming public. Such work should proceed with increased energy as experience indicates what are the most effective methods. Apple advertisers now have an increasing amount of authority for proclaiming the unusual health-giving qualities possessed by the apple, and failure to capitalize upon the health appeal would certainly be a serious mistake.

Fruit growers should think this matter through carefully, and support to the fullest extent the work of the apple institutes. The citrus growers have done a commendable piece of national advertising, and, as a result, have been able to market increasing supplies of citrus fruits every year. Apple growers can well afford to follow suit and should deliberately set about not only to hold their present domestic market, but to expand it. Even though better times for the fruit grower seem just ahead, no effort should be spared to maintain the gains already achieved through the instrumentalities already at work.

ANALYSIS OF ORCHARD MANAGEMENT

PROF. A. H. TESKE, extension horticulturist, Blacksburg, Va., presented a clear-cut analysis of "The Orchard Management Program for 1937" at the annual convention at Roanoke. This thoughtful paper dealt generally and specifically with many angles of fruit growing. Some pertinent paragraphs are herewith presented because of their timely value:

With a reduction in the number of apple trees and a decrease in both total and commercial production, and with greater stability in the industry, the apple grower today still finds it difficult to market this reduced supply at satisfactory prices except in years when there is an unusually short crop such as we had this past year. Growers may well inquire as to the reasons. Why is it so hard to dispose of their fruit at fair prices in such crop years as we had in 1935?

A part of the answer can be found in the fact that up to the time of the depression more than 16 per cent of the commercial apple crop of the United States and from 40 to 60 per cent of the commercial apple crop of Virginia went into foreign markets.

Again, while we have been pulling out apple trees by the millions, other countries have been increasing their plantings so that, although apple production in the United States is showing a downward trend, world production is on the increase.

Another reason why apple growers are having trouble in marketing medium to

large crops at satisfactory prices is that production of competing fruits is on the increase. This is especially true of oranges and grapefruit. The production of these fruits is already so large that during the past three crop years considerable quantities of oranges have been withheld from the fresh fruit markets because they would not pay handling charges. Increased quantities of grapefruit have also been going to the canneries each year. Prices of citrus fruits are expected to go lower as production increases.

As prices of oranges and grapefruit dip to lower levels, it can be expected that the citrus industry will launch a more vigorous campaign to increase the consumption of these fruits. All of which means that in the future the apple industry must be prepared to meet stronger competition.

With prospects for an increase in the world's supply of apples and other fruits, with foreign markets lost to a large extent and little hope of a return to the good old days of a free and open door, and with stronger competition from increased citrus supplies for the consumer's dollar in the domestic markets, the situation in which the apple industry finds itself is entirely different from that of the late '20's. It calls for a further adjustment in the business of fruit growing.

I believe that most apple growers will agree that it takes a well fed, vigorous and healthy tree to produce a good crop of quality fruit, and that we cannot expect

much from broken down, weak and sick trees. And is it not true that the apples that usually come from such trees lack size and color? And is it not the fruit from such trees that increases the amount of low-grade fruit which, when dumped on the markets, slows up the apple movement and lowers the prices of the better grades? Why spend time and money on pruning, spray materials and fertilizers to coax these poor trees along, when, as a general rule, they lose money for the grower and make his marketing problems more difficult?

As the first step in the management program for the coming year, I would suggest taking out such trees. This would also apply to trees that are planted so close that there is excessive crowding at the expense of satisfactory yields.

With the weak, sick and partly dead trees out of the way, the next question that should be of immediate interest to the apple grower at this particular time is that of fertilization. Growers should bear in mind that we had a short crop in 1936, and that there is a heavy set of fruit buds for the 1937 crop. If there is a big crop in 1937, prices are likely to be lower.

Some growers have already made up their minds not to use any fertilizer on their trees this year. Many have asked the question as to what would be the best fertilizer to use.

This is one of the years when the grower cannot afford to omit the fertilizer application to his trees. Best results cannot be secured from a hit and miss fertilizing program. The only thing apple growers have to sell is the raw materials of the soil, manufactured into plant foods by the leaves of their trees and transformed into a usable product, in this case apples, for which consumers are willing to pay a price. If apple trees are to function properly and efficiently, they must be supplied with the materials which go to make for performance. They must be fed. And, within limits, the more raw materials that are put through the factory—that is, the leaf surface of the apple trees—the larger will be the output of that factory.

We know that beginning with the swelling of the buds up through the blooming and fruit setting period, apple trees require large amounts of nitrogen. And, at that early season period, nitrates are less plentiful in the soil. In addition to the nitrogen requirements for blooming and fruit setting, the trees need a good supply of nitrogen to develop a good dark green leaf surface capable of manufacturing food later for the development of fruit and the formation of fruit buds.

During the 1936 short crop year, the apple trees formed a large number of fruit buds. This means that this heavy set of flower buds will use up much of the supply of reserve food in blooming this coming spring. The competition for the reserve food between the heavy set of flowers and the first leaves will be so strong that the early leaf surface is likely to be so small that it cannot manufacture enough starch foods to meet the early season's growth requirements and at the same time provide for a surplus to form fruit buds for the next year. This means that unless we do something to increase the reserve food supply, there will be only a small number of fruit buds formed for a crop in 1938.

In my opinion, soil erosion and the lack of organic matter in our orchard soils are the two most important problems of the apple industry at the present time. The grower whose orchard management program is a satisfactory answer to these problems and nitrogen requirements of his trees will have gone a long way toward solving the problem of increased yields.

H. L. Lantz
SECRETARY

APRIL, 1937



"BLACK LEAF 40"

Kills Aphis and Other Insects

BLOSSOM TIME is promise time—foretelling the possibilities of your crop. But if these promises are to be fulfilled, you must aid in guarding against attacks by Aphis, Red Bug, Leaf Hopper and other insects which damage foliage and cause dwarfed or gnarled fruit. "Black Leaf 40" used alone or with other standard sprays kills these pests—both by contact and by fumes.

FOR MORE EFFECTIVE CODLING MOTH SPRAYS: Fortify lead arsenate or "summer-oil" sprays with "Black Leaf 40". Lead arsenate kills the larvae after they eat, "summer-oil" kills the eggs. "Black Leaf 40" kills mature eggs and larvae, and also adult moths.

"BLACK LEAF 40" IS EASY TO USE: Concentrated, effective, easy to mix and apply. Of vegetable origin—it is not caustic—does not "burn" man, horses, trees or crops. Volatile, it "fumes off" (evaporates) from foliage or fruit. "Black Leaf 40" is sold by spray material dealers everywhere.

3733

TOBACCO BY-PRODUCTS AND CHEMICAL CORP. INCORPORATED
LOUISVILLE KENTUCKY

STATE NEWS

FROM NEAR AND FAR



ILLINOIS—Fruit crop outlook for Illinois: Promising, particularly for apples, peaches and pears. Anticipated 1937 crop below crop of 1935, however, due to severely cold winter of 1935-36 followed by intense heat and drought of last summer which killed a large number of apple and peach trees.

Summer apple territory in Johnson, Union and Jackson counties will have only a reasonable crop, many orchards having a very light bud set.

Peaches in the territory south of Carbondale were severely injured during the winter and only a partial crop is expected.

Strawberries do not promise a heavy yield in any section of the State.

Peaches, apples and pears in Centralia area have encouraging outlook. Peaches in most orchards have survived the winter in sufficient numbers to promise a full crop.

Western and northern Illinois are due a crop of average production.

Present indications: Illinois will have a fruit crop distributed over entire producing area, perhaps about 80 per cent of 1935 production.—**JOE B. HALE**, Sec'y, Salem.

TENNESSEE—Poisoning, burning, electrocuting, trapping, mashing, and starvation are some of the methods used to destroy codling moths and their larvae.

N. N. Barnett of Troy, Tenn., tried out another common means of ending life—



drowning. He found a number of worms spun up in a leather and canvas backband, which had been discarded in his orchard. He threw it into a barrel of rainwater and after three days under water, found all the worms alive. He left it in the barrel for two weeks more and reports that many of the worms were still alive.

As far as we know, Mr. Barnett is not a member of the Tall Story Club.

On Lincoln's Birthday, 27 members of the Greenfield (Weakley County) Fruit Growers Association gathered for their annual meeting. Vice-president R. R. McUmber told the writer this association has operated for 40 years under but three presidents. They employ a sales manager, and one man has held this position for 20 years. We believe this remarkable record reveals the fine character of its members.

Denton Fly, Milan, 1936 champion strawberry grower, won further honors and a \$500 cash award, given by the *Commercial Appeal*, Memphis newspaper, for the best all around farmer in Tennessee, Arkansas and Mississippi area covered by its circulation.

Purpose of the award was to encourage diversification of crops and required keeping of account records. Mr. Fly's records showed a profit on 14 out of 16 crop projects.—**A. N. PRATT**, State horticulturist, Nashville.

MINNESOTA—Grower reports indicate most orchard trees and small fruit plants appear to be coming through the winter in good condition.

Although the area around Minneapolis and St. Paul is still very deficient in moisture, un-

PAGE 20

usually heavy snowfall with comparatively little frost in the ground in the rest of the State gives promise of favorable planting conditions for the spring of 1937.—**J. D. WINTER**, Sec'y, St. Paul.

CALIFORNIA—Enterprising San Joaquin Valley farmers will merely grin when you tell them, "You can't eat cactus!" For they are now growing a cactus crop that makes real money for them. The product is prickly pears.



Each row of cactus reaching across a 30-acre field, for instance, pays \$250 to \$300 annually.

The pears are grown at the end of the big heavy cactus leaves, and a part of the latter is cut off when harvesting. This practice, it has been found, keeps the pear in good condition for a month or two.

The fruit is eaten sliced, with sugar and cream.

One American authority says Americans contract some 100,000,000 colds every winter. Capitalizing on such catastrophe, California Fruit Growers Exchange conducted an intensive seasonal campaign to sell lemons for "Hot Lemonade for Colds."

SOUTH DAKOTA—The 32 seedling apricots which Dr. N. E. Hansen of S. D. State College collected in 1924 from Old Man Winter's headquarters in eastern Siberia—where the temperature ranges from 47 to 50 degrees below zero—have been bearing for several years. Dr. Hansen feels these new apricots, the fruit of which is yellow, somewhat smaller than the apricots of commerce, and makes delicious preserves, will stand the extreme conditions of the Dakotas, Minnesota, Montana, Manitoba and Saskatchewan.

PENNSYLVANIA—The morbid facts back of the serious winter injury suffered by Pennsylvania fruit trees in 1933-34 and 1935-36 were disclosed by Dr. R. D. Anthony of Pennsylvania State College at the annual meeting of the hort association. Among the factors were in-



adequate soil drainage, soil fertility, and heavy bearing. Dr. Anthony bases his facts on a tri-state survey made the past season.

Recently a number of cases have been observed where the heavy drain of a large crop and its effect upon the degree of winter injury is evident. In a block of Stayman trees carrying a heavy bloom in 1935 a satisfactory crop was produced except upon two rows of trees that were not covered during the critical scab period due to sprayer trouble.

The two rows not fruiting in 1935 showed no winter injury in 1936 and produced a maximum crop whereas the remainder of the trees carried a very light crop and suffered severe injury.

AMERICAN FRUIT GROWER

A second orchard of Baldwins carried a heavy crop in 1935. The fruit on two and a half rows of trees was thinned the latter part of June. In 1936 the trees upon which the crop was limited in 1935 by thinning appeared to be normal while the unthinned trees were seriously damaged, about 50 per cent passing out of production.

A third case is that of an orchard comprised of Delicious and York Imperial receiving heavy applications of nitrate. Here again the trees carrying heavy crops in 1935 were damaged to such an extent that removal is the only course to follow.

In the past the practice of fruit thinning was emphasized as a means of improving the quality of the pack but in some Pennsylvania orchards conditions would indicate that fruit thinning might play a more important role in maintaining tree vigor and long-time production.—**J. U. RUEF**, Sec'y, State College.

(Readers are referred to Dr. Anthony's and Dr. R. H. Sudds' article, "Low Temperature Injury to Orchards," January issue.—Editor)

WISCONSIN—All northern states patiently await the outcome of the tests being conducted with Crath Carpathian English walnuts (introduced from Poland), seed of which are again being distributed this spring for trial by Wisconsin Horticultural Society.



So far the tests have been successful. Seedling trees have come through the winter in good condition. Older trees have survived our severest winters. None of the trees in the U.S., however, have borne any nuts as far as we can determine. Trees in this country should produce at about seven or eight years.

Interest in new fruit varieties is increasing. Many older varieties have been discarded for the new. Wisconsin growers are becoming more favorably impressed with Cortland and Kendall to lengthen the season of McIntosh, the State's favorite apple.

Melba has become a standard early variety for Wisconsin. Beacon is replacing Duchess in Minnesota and is being widely tested in Wisconsin this year.

It is indicated that Haralson will probably replace N.W. Greening as a cooking apple and be planted extensively in sections having severe winters.

A number of Wisconsin Hort Society members are testing the new hardy apricots introduced from China by Dr. N. E. Hansen of South Dakota.

In the strawberry world Beaver and Premier continue to retain their popularity in commercial sections, although hundreds of growers are experimenting with Dorsett, Fairfax, and Catskill. These varieties look very promising and will probably replace other varieties on soils and under conditions where they can be most successfully grown.

Rather surprising differences in results are being noted between different varieties depending on soil conditions and locality. For example, Beaver has been a failure in many sections of the U.S. but continues to be a

(Continued on page 22)

APRIL, 1937

GEORGIA PEACH ORCHARD'S COVER CROP

turned under on time!



While the fertilizing value is greatest—and before it wastes moisture needed by the coming peach crop—this rich cover crop of Austrian Peas is turned into the soil.

No worry over traction—the Twenty-Two's broad tracks keep their sure footing even in damp, slippery cover crop. No time lost working over grades or side-hills—this rugged tractor is balanced to keep its tracks on the soil their full length. And the full-pressure-lubricated engine develops heavy-duty power to team with non-slip traction.


With the 6-foot, offset cover crop disk, this "Caterpillar" Twenty-Two Tractor thoroughly cultivates the rolling 85-acre peach orchard

in only five 10-hour days. And using only 2 gallons of low-cost fuel per hour!

Besides its orchard jobs, this

tractor also helps Owner Lamar Potts, Hogansville, Georgia, raise 400 acres of cotton, corn and oats!

CATERPILLAR TRACTOR CO. PEORIA, ILL.



REG. U. S. PAT. OFF.

WORLD'S LARGEST MANUFACTURER OF DIESEL
ENGINES AND TRACK-TYPE TRACTORS

CATERPILLAR TRACTOR CO.,
Dept. A-4, Peoria, Illinois

Gentlemen: I operate _____ acres of orchard.

My present power is _____

Please send information on:

☐ Orchard Model Twenty-Two

☐ Orchard Model Thirty

☐ Orchard Model Diesel RD 4

☐ "Caterpillar" Power for Orchards


Name _____ R. F. D. _____

Town _____ County _____ State _____


Camera!



H. L. Lantz, left, secretary of the American Pomological Society, pauses to chat with W. S. Campfield, center, secretary of the Virginia Horticultural Society, and G. Leslie Smith, president of Illinois Horticultural Society, at the Virginia meeting.



Left to right: Francis G. Reiter, fruit grower of Mars, Pa.; J. U. Ruef, Pennsylvania State College extension horticulturist and new Pennsylvania association secretary; George A. Goodling, Loganville; and C. B. Snyder, Ephrata, Pennsylvania, Horticultural Association treasurer, look directly at the American Fruit Grower camera for this "shot" at the annual meeting of the association.



H. W. Miller, left, veteran West Virginia fruit grower, and J. B. Whisnant, right, prominent Georgia fruit producer, look over the program of the Virginia meeting with Dr. H. P. Gould of the U.S.D.A.

STATE NEWS

(Continued from page 20)

very profitable variety in certain sections of Wisconsin. Hence we urge growers to try different varieties and adopt those best suited to their soil.

Fruit growers are being asked to co-operate with beekeepers in an effort to obtain a large appropriation to drive foul brood (disease of bees) from the State.—H. J. RAHMLow, Sec'y, Madison.

INDIANA—Practically all small growers of apples have been sold out for many weeks and larger growers had comparatively few apples still to be moved out of storage as we entered the last half of March.

Indications at that time pointed to a very heavy apple bloom throughout the State.

Moisture conditions in orchards are tops. Even with less-than-average rainfall the next few months, deeper soils are expected to carry an adequate moisture supply farther into the season than in 1936.

Considerably fewer orchards are operated under clean cultivation today as compared with a few years ago. Erosion problems, moisture-holding capacity, soil depths and soil temperatures have all contributed toward this gradual change of soil management throughout the State.

Free to Hoosiers is 36-page Bulletin No. 414, edited by Clarence E. Baker and entitled, "The Relation of Nitrogen and Soil Moisture to Growth and Fruitfulness of Apple Trees Under Different Systems of Soil Management."

Comparisons are given of commercial apple production on clean cultivated plots as compared with blue grass and alfalfa-sweet clover sods, both with and without commercial fertilizer applications.

Purdue Agricultural Experiment Station, West Lafayette, will send the bulletin upon request.—EVERETT WRIGHT, Sec'y, Lafayette.

MASSACHUSETTS—The 43rd annual meet of Massachusetts Fruit Growers Association in Worcester was warmed up with a jolly attendance of over 500 fruitmen. Convention was held in co-operation with the Union Agricultural meetings sponsored by the State Department of Agriculture.

Discussions on orchard heating brought out the fact that heat is more important than smudge and that fruitmen should experiment with their particular situation before they go to too great expense in trying to prevent frosts in their orchards.

Fruitmen who have tried orchard heating stated further it was necessary to keep constant watch on the temperature and to have everything set for lighting the fires when the temperature got down to 31 degrees F, that being the point at which fires should be lighted.

The association went on record as objecting to renaming of apple varieties to avoid confusion. Standardization of varieties and variety names certainly should be maintained. If we were to add, say, one name a year to a variety, by 1950 the situation would be comic.

McIntosh is still New England's No. 1 apple, according to Dr. Richard Wellington, Geneva (N.Y.) Experiment Station.


Shining light of the fruit show: Known Orchards of Sutton. This orchard took both sweepstakes prizes, one on the bushel package class and one on the plate class.—WILLIAM R. COLE, Sec'y, Amherst.

WASHINGTON—A bill providing for a tax of two cents per cwt. on apples to support a national advertising campaign was recently passed by Washington legislators.

This is the same assessment as has been paid by members of Washington State Apples, Inc., growers' organization which launched industrial advertising on Washington apples on a voluntary basis this season.


It is expected the growers' organization will be taken over by the State to administer the new program.

Farmers are saying
"SEE GMC FOR EXTRA VALUE"



IN THE farming industry—in every industry—truck buyers, with their greater profit in mind, are looking to GMC for extra value. Whatever your hauling needs may be, you'll find in the extensive GMC line a "truck-built" vehicle whose ability to perform ably and save money has already been conclusively proved. In the half-ton range, there are GMC's in both 112-inch and 126-inch wheelbases and with the largest standard bodies in the industry. In the 1½-2 ton range, there is a GMC in several wheelbases and with

features that set it apart as an unmatched value. And, of vital importance, every GMC has advanced stream-style and exclusive "dual-tone" color design that assure an exceptionally distinctive appearance. Get the facts! See GMC for extra value!



**QUALITY AT PRICES
 LOWER THAN AVERAGE**

Our own Y. M. A. C. Time Payment Plan assures you of lowest available rates

GENERAL MOTORS TRUCKS & TRAILERS

GENERAL MOTORS TRUCK & COACH

DIVISION OF

YELLOW TRUCK & COACH MANUFACTURING COMPANY, PONTIAC, MICHIGAN

AMERICAN FRUIT GROWER

APRIL, 1937

PAGE 23




DOING THE WORK OF SIX MEN—

The MOTOR SCYTHE



PAYS FOR ITSELF in short order

● For mowing between and under low trees the Motor Scythe can't be beat. It reaches in-under where it is almost impossible to mow with a hand scythe. Sturdily built, perfectly balanced, easily operated, 30 inch wheel, 36 inch cutting bar mows 30 inch swath. 3/4-horse power motor operates 4 hours on 1/2 gallon of gasoline. Weight, 135 pounds. List price, \$125.00 (F. O. B., Detroit). For full particulars address:



DETROIT
HARVESTER
COMPANY

5457 W. Jefferson
Ave.,
Detroit, Mich.

**Detroit Harvester
MOTOR SCYTHE**



Genuinely
Friendly

featuring—

Unusually Comfortable Rooms,
Good Food, Faultlessly Served and
Uniform Rates, starting at \$2.75
single.

In CLEVELAND it's

● **THE HOLLENDEN**
RICHARD F. MARSH, Mgr.

In COLUMBUS it's

● **THE NEIL HOUSE**
TOM. A. SABREY, Mgr.

In AKRON it's

● **THE MAYFLOWER**
C. J. FITZPATRICK, Mgr.

In TOLEDO it's

● **THE NEW SECOR**
B. R. BRENAN, JR., Mgr.

For Your Winter Vacation

In MIAMI BEACH it's

● **THE FLEETWOOD**
JOHN HEARST, Mgr.



Aerial view of Florida experimental vineyards. Plots appearing as roadways are actually defoliated check plants.

GRAPES

FLORIDA TESTS

Striking results of a grape spraying experiment are shown in the above illustration. This work was conducted by Kenneth W. Loucks at the Leesburg Field Station of the Florida Agricultural Experiment Station. Of the many materials tested for the control of the grape rots, Bordeaux mixture proved best. The tests were conducted over a period of years and the results have proved of untold value to Florida grape growers.

BERRY MOTH CONTROL

Entomologists of the New York State Agricultural Experiment Station at Geneva have warned grape growers to be on the lookout for signs of the grape berry moth in their vineyards. Periods of severe cold weather when the ground is not covered with snow usually account for the destruction of large numbers of the overwintering larvae of the berry moth, say the specialists, but in some vineyards protection by snow may have allowed a sufficient survival to make possible troublesome infestations. Grape growers are urged, therefore, to take the necessary measures to combat this pest before it is too late.

"Examine the blossom clusters frequently until the grapes are the size of small peas, for any signs of webbing by the larvae of the grape berry moth," says Prof. F. Z. Hartzell, New York station entomologist. "The young worms spin webs in the clusters which are usually readily detected. If the insects are present in any number, a thorough application of spray should be made immediately after the grapes have set, or at least by the time they have attained the size of buckshot.

"Every effort should be made to eliminate the pest just after blossoming, as later applications will intro-

duce spray residue problems. Efficient control can be secured by a thorough spraying of the young berries."

Experiments in the eastern grape sections have shown that the berry moth can be controlled by spraying with arsenate of lead at the rate of three pounds to 50 gallons of Bordeaux mixture. One application of this spray is made shortly after fruit setting, another about 10 days later, and the last when the grapes are about half grown.

EARLY FRUIT BEARING

A method whereby seedling grapes of California varieties may be brought into bearing in 18 months, instead of the usual four or five years, has been developed by U.S.D.A. scientists as a means of speeding up experimental breeding work with this fruit.

At the Fresno, Calif., experiment station, Elmer Snyder and F. N. Harmon of the Bureau of Plant Industry have used the simple "T" bud as a means of hastening fruit bearing of promising new varieties.

The seeds are planted about the first of February in a greenhouse flat. A few weeks later, when the plant has come up and a few leaves have started to develop, it is transferred to a gallon can or a six to eight-inch greenhouse pot. By the latter part of May or first part of June the seedling attains a growth of 12 to 16 inches, from which three or four buds can be obtained.

These are "T" budded into vigorous rootstocks or bearing vines in the vineyard. They are wrapped securely with rubber grafting tape and the cane is pinched off at the tip to check terminal growth. A week to 10 days later the shoot is cut back to the bud, leaving some of the original leaves as shade. As three or four buds are inserted in each shoot, the weaker ones are pinched back and only the strongest allowed to grow. These shoots, under the conditions at Fresno, will bear fruit in August of the second year, about 18 months after the original seeds were planted.

During the past year Snyder and

Harmon made over 1,500 "T" buds and about 80 per cent grew successfully. The buds must be shaded during extremely warm weather the first summer. It is also important to cut the shoots back at the proper time, and constant upright training was found to promote more rapid growth.

Although final commercial merits of a promising seedling cannot be decided this early, many qualities, such as color, shape, size and flavor, can be determined at this time, thereby speeding experimental work.

PLUM PRODUCTION

(Continued from page 14)

but is a good annual producer and starts to bear at an early age.

Yellow Egg is well worth mentioning as it is very attractive, good for the roadside market, and has quality that many of the other even less acid ones do not have.

The purple Bradshaw attains good size, is good for eating fresh when ripe, and sells well at roadside stands and to hucksters. Early to ripen, it starts off the plum market. The tree is large and long lived.

I have described varieties I consider most desirable to plant, stressing those in the prune class as my experience and observation convince me these are the superior fruits.

So far as diseases and insects affecting plums are concerned, these are as easily controlled as those of the apple. The trees are subject to scale and occasionally to red mite. Curculio is the worst insect enemy, but sprays applied properly will control it. Leaf spot disease, to which they are susceptible, can be controlled by combining a fungicide with the insecticide.


I began thinning plums years ago, endeavoring to prevent contact of specimens. The clusters hold moisture during wet times and encourage rots. Thinning of plums, however, is not as generally practiced as with larger fruits.

Pruning is done by detailed thinning of the fruiting surface—not by cutting out large branches.

I usually cultivate until midsummer, then sow a cover crop where the trees do not make too much shade, or let the planting grow to native grasses and weeds. Most of the soil is tiled for good drainage. Some of my trees are on sandy soil, but I have found that this type of soil is not as good as heavier land for plums. For fertilizers I have used nitrate of soda, cyanamid, sulphate of ammonia, and manure at different times.

Improved varieties, along with proper methods of pest control and soil management, must be considered by the plum grower if he is to obtain the greatest benefit from his enterprise.

"FORD'S GOT THE RIGHT IDEA"



No matter what crops you raise . . . no matter what your farm hauling problem may be . . . there's a Ford V-8 Truck or Commercial Car that will handle your job and do it at low cost. That's why farmers this year are changing to Fords and finding a new way to make more net profit on their crops.

Every type in the Ford V-8 Truck and Commercial Car line is now available with an 85-horsepower V-8 engine improved to give greater economy in heavy-duty and high-speed service. In addition, those types most often used for light loads are available with the new 60-horsepower V-8. This new engine has already proved itself a real money saver to the farmer whose loads are light.

Get the facts about V-8 economy. Find out for yourself how a Ford V-8 Truck or Commercial Car can cut your farm hauling costs. Ask your nearest Ford dealer for an "on-the-job" test under your own operating conditions.

● Convenient, economical terms through Authorized Ford Finance Plans of the Universal Credit Company



FORD V-8 TRUCKS AND COMMERCIAL CARS

RENEW YOUR SUBSCRIPTION NOW!

SAVE Money—Order at
Low Cost Your Favorite
Magazines

Club No. 101
AMERICAN FRUIT GROWER.....1 yr. } ALL
Woman's World.....1 yr. } SIX
Good Stories.....1 yr. } ONLY
Home Circle.....1 yr. } \$1.00
Illustrated Mechanics.....1 yr.
Mother's Home Life.....1 yr.

Club No. 105
AMERICAN FRUIT GROWER.....1 yr. } ALL
Pictorial Review.....1 yr. } FIVE
Pathfinder (weekly).....1 yr. } ONLY
Good Stories.....1 yr. } \$1.60
Illustrated Mechanics.....1 yr.

Club No. 111
AMERICAN FRUIT GROWER.....1 yr. } ALL
McCall's Magazine.....1 yr. } FOUR
Household Magazine.....1 yr. } ONLY
Illustrated Mechanics.....1 yr. } \$1.40

Club No. 112
AMERICAN FRUIT GROWER.....1 yr. } ALL
Pictorial Review.....1 yr. } FOUR
Woman's World.....1 yr. } ONLY
Household Magazine.....1 yr. } \$1.45

Club No. 114
AMERICAN FRUIT GROWER.....1 yr. } ALL
Breeder's Gazette.....1 yr. } FIVE
Woman's World.....1 yr. } ONLY
Good Stories.....1 yr. } \$1.00
Poultry Tribune.....1 yr.

AMERICAN FRUIT GROWER
For One (1) Year and Your
Choice of Any Three of the
Following Magazines for
ONLY \$1.00

Check Three
() American Poultry Journal.....1 yr.
() Cloverleaf American Review.....1 yr.
() Everybodys Poultry Magazine.....1 yr.
() Gentlewoman Magazine.....1 yr.
() Rhode Island Red Journal.....1 yr.
() Successful Farming.....1 yr.
() Michigan Farmer.....1 yr.
() Pathfinder (weekly).....26 issues
() Illustrated Mechanics.....1 yr.
() Plymouth Rock Monthly.....1 yr.
() Mother's Home Life.....1 yr.
() Good Stories.....1 yr.
() Home Circle.....1 yr.
() Ohio Farmer.....1 yr.
() Household Magazine.....1 yr.
() Woman's World.....1 yr.
() Leghorn World.....1 yr.
() Poultry Tribune.....1 yr.
() Capper's Farmer.....1 yr.
() Breeder's Gazette.....1 yr.
() Home Arts Needlecraft.....1 yr.


AMERICAN FRUIT GROWER, 1370 Ontario St.,
Cleveland, Ohio.

Enclosed find \$..... for which please send me
the magazines marked with an X.

Name

Postoffice

R.F.D. State



In drying apricots, the fruit is halved, pit removed, placed in trays with the cups up, and exposed in a tight compartment to burning sulphur fumes. After removal from the sulphur house, the trays are spread on the ground and exposed to the sun for one to three days. The trays are then stacked, as shown in the background of above photograph, and drying completed in the shade, which takes another six or seven days. Drying ratio of fresh to dry fruit is about five to one.

APRICOT CULTURE

(Continued from page 13)

in California consider the apricot as probably the most difficult tree and crop to handle. Those who have grown both apricots and sweet cherries are somewhat divided in their opinion as to which of these two fruits requires the greater skill on the part of the orchardist, with the majority perhaps voting for the culture of the apricot as presenting the greatest number of hazards.

The tree itself in most locations presents no great problem, being grown with almost equal success on apricot or peach root, and being adapted to heavier and more poorly drained soils by working on Myrobalan, certain strains of which are fairly satisfactory as a rootstock. The largest apricot trees that I know of in California are on peach roots and are now about 80 years of age. Figure 1 shows one of these trees with a spread of more than 60 feet. This tree has several times produced more than a ton of fruit in a season. Many orchards in California are over 50 years of age and apparently have many more years of profitable fruiting ahead.

Although the trees are apparently thrifty, and long lived even when not grown under ideal conditions, the production of a profitable annual crop may be an altogether different matter.

Fruit buds of the apricot are apparently as hardy to low winter temperatures as are those of the peach, but the blossoms and young fruits are the most susceptible of all deciduous fruits to a few degrees of frost, and in fact at certain stages of development a temperature of 31 1/2 degrees F if continued for an hour or two will cause loss.

During the dormant period tem-

peratures must not be too mild or else the "rest" will not be broken. After unusually warm winters many fruit buds drop rather than opening, thus resulting in a decided reduction in crop. Where irrigation facilities are available, the apricot grows successfully under climates ranging from cool coastal districts to desert conditions. The time of harvest is dependent upon the accumulation of a certain number of heat units, whether secured in 100 or 130 days, a fact explaining the earliness of certain districts. Temperatures of about 103 degrees F and upward during the ripening season cause "pit burn"—a darkening of the flesh around the pit—which spoils the fruit for fresh consumption or canning and lowers its value for drying.

Adequate soil moisture must be present at all times if regular and profitable fruiting is to be obtained. Irrigation facilities are necessary to maintain sufficient new growth and annual crops. From two to eight irrigations per season are given depending upon the district and soil type. Some varieties, such as the Tilton, seem to be less tolerant of drought than others, such as the Royal.

On account of the ultimate size of the apricot tree under suitable soil conditions, it is questionable whether they should ever be planted closer than 24 by 24 feet. All varieties grown are apparently self-fertile and no provision for cross-pollination is necessary.

One-year-old trees are usually planted and headed to a whip at about 30 inches above the ground. Three main branches are selected at the first winter pruning and generally

headed at about shoulder height. At the second annual pruning from five to seven secondary scaffolds are saved without heading. Thereafter, the trees are "thinned" each year so long as adequate replacement wood can be secured without recourse to "heading." Fruit is produced on one-year shoots and short-lived spurs. After the peach, the apricot probably receives the most severe pruning of the various deciduous tree fruits grown in California. In addition to limiting the crop by pruning, hand thinning of the fruit is almost universally practiced.

Bacterial gummosis, brown rot, black heart, and shot hole fungus (*Coryneum*) are the most serious diseases. Insect pests include the peach root borer, peach twig borer, canker worm, leaf roller, brown apricot scale, and black scale. Except bacterial gummosis, black heart, and the root borer, all these troubles may be controlled by known spray programs. Amputation and scarification, plus a disinfectant, afford some control of bacterial gummosis. Paradichlorobenzene has proved useful against the root and crown borer.

The Royal, an old French variety, and the English variety, Blenheim (Shipley), are most widely planted. These two sorts were undoubtedly distinct varieties as first introduced but have become decidedly confused in more recent years, so that few, if any, can with certainty distinguish between them. Both of these varieties are entirely satisfactory for fresh consumption, canning, and drying.

During the past 30 years a California seedling, the Tilton, has been planted rather widely in the interior valleys. It is larger than either Royal or Blenheim and produces heavy crops under ideal conditions. Its quality for either drying or canning does not equal Royal or Blenheim.

The Moorpark, another old English variety, is of the highest quality for all purposes but has almost passed out of cultivation on account of shyness in production. The so-called Wenatchee Moorpark is popular in Washington and some of the other western states.

The Derby, Stewart, Newcastle, and Wigan are occasionally planted in early districts for fresh shipment. The latter two are rather soft and must be harvested too firm for best quality. The Derby and Stewart closely resemble the Royal and are often shipped as such.

Apricots are generally picked by hand, proper maturity being determined largely by color. For distant shipment a full straw color and complete freedom of flesh from the pit is the maturity criterion; for the cannery and dry yard, the fruit is left on the trees until much softer, being most mature for drying. After picking there is no increase in sugar content.

In California, apricots for the east-

(Continued on page 35)

"1/2 GALLON OF GAS PER ACRE"

Cultivates 26 acres of corn in 8 hours with a high compression Oliver "70"



"With high compression, good gasoline is more economical than ever," says Merritt Klopfenstein of Dundee, Michigan, who farms 260 acres. Mrs. Klopfenstein runs the Sunny Ridge Hatchery, with a 40,000-chick capacity, making a specialty of ducklings and turkey poults.



THE economy of good gasoline is an old story to Merritt Klopfenstein, Dundee, Michigan. He's been using it more than ten years in his old low compression tractor, and has only had the valves ground twice and no major repairs. But the performance of his new high compression Oliver "70" opened his eyes.

He writes, "I've done my first full season's work with my high compression Oliver '70' and it's more economical, more powerful, and covers the ground faster. For example, I cultivated 26 acres of corn from 7 o'clock at night to 3 o'clock in the morning, and in eight hours steady work, used only one-half a gallon of gasoline to the acre.

"When I used my new Oliver and

my old tractor together, the Oliver made four rounds of the field to the other's three. Our first high compression tractor has proved to us that good gasoline is more economical than ever, and that is why we are going to trade our old low compression tractor for another new high compression Oliver Rowcrop '70.'"

Good gasoline in your car, truck or tractor steps up the useful power, gets work done faster, eliminates costly repairs, and in your tractor, reduces oil bills by as much as two-thirds. Get this added economy, performance, and speed by using regular grade gasoline.

Ethyl Gasoline Corporation, Chrysler Building, New York, manufacturers of anti-knock fluids for premium and regular gasolines.

It pays to buy **GOOD GASOLINE**
FOR CARS, TRUCKS AND TRACTORS

Why

it costs less to spray with a **HARDIE**

Vertical crankshaft driven pump with long connecting rods avoiding side-thrust.

All important parts are separate units easily accessible and economically replaced.

Volumetric pump efficiency 96% to 99%—highest in the industry.

Delivers full rated capacity and pressure easily all the time at slow speed.

Individual selective lubrication with clean oil for every moving part.

Fully lubricated plunger and cup. A feature not found in any other pump.




Roller bearing trucks. Interchangeable steel or wood tanks.



● Hardie sprayers are designed and built to cut the cost of spraying. The vertical crankshaft driven pump; the complete selective lubrication of every moving part *including plunger and plunger cups*; the ready accessibility of every part making for quick, easy, economical service; the complete dust-proofing of pump and engine; the roller bearing trucks—each and every part of a Hardie sprayer specifically and directly reduces the cost of spray application. ● Then consider the long, long life of the Hardie—the years of use in which the owner gets rich dividends on his investment—no trouble, no loss of time nor material, practically no upkeep expense. Whatever may be your spraying requirements—large or small—insist upon having Hardie efficiency and economy.

THE HARDIE MFG. COMPANY HUDSON, MICH.

Branch Factories, Sales and Service Offices:
Portland, Ore., Los Angeles, Calif., Kansas
City, Mo., Brockport, N. Y., Hagerstown, Md.
Export Dept., Detroit, Mich.



You must have a 1937 Hardie catalog to learn about the really new things in the sprayer industry. We will gladly send you a copy of this beautiful 64-page book. Write for it.

HARDIE

DEPENDABLE SPRAYERS


CONTROL OF CHERRY AND PLUM BLACK KNOT

By M. B. CUMMINGS
University of Vermont


BLACK KNOT disease is not new but is a very troublesome and recurring one. It often appears in great abundance on uncultivated plums and cherries, and spreads to cultivated ones.

The accompanying illustration shows a tree that is heavily infected with black knot. This is an extreme case and a tree in this condition should receive immediate attention. In this instance, it would not be worth while to try to save the tree by cutting out affected limbs or individual knots. The tree is too much knotted. It would be better to cut the tree off at the ground and put the whole specimen on the bonfire.

But many times one may find only a few knots in a plum or cherry orchard or on trees in the fence row or in the garden. Such trees can be cleared of black knots and the trouble prevented from spreading to other cultivated trees. If this work is done before the season opens and before the spores have spread, good control should be secured. Cut the limbs one foot below the knot so as to remove all infected tissue. Black knot extends down the tissue farther than is apparent on the outside.



Black knot disease of plum with swellings all over the tree. If the knot is cut off and burned before the season opens, spread of the trouble is forestalled.



Peach thinning, as practiced by these workers, is an important part of every peach grower's cultural program.

PEACH THINNING

(Continued from page 11)

The first extends up to about the time the pit begins to harden at the tip. During that time the stone reaches its full size, although the flesh is as yet quite thin about it. During the second growth period the enlargement in the peach as a whole is relatively slight. This lasts in Elberta up until five

is reduced on overloaded trees. Since fruit bud formation is so intimately connected with growth, unthinned trees have sometimes produced no more than 10 per cent as many fruit buds as the thinned trees. The color of the fruit is also increased by thinning, and the time of ripening is

TABLE I. The relation between thinning, nitrating, and pruning in controlling size. Elberta season of 1931.

Orchard	Treatment			Age of trees	No. of trees	Yield of fruit in pounds of the different sizes					Av. yield per tree (lbs.)
	Nitrogen	Pruning	Thinning			0-1 3/4"	1 3/4"-2"	2-2 1/4"	2 1/4"-2 3/4"	2 3/4"-up	
1. C. F. Heaton New Burnside	3 lbs. Sulphate of ammonia	Regular orchard mod. heavy	None	13	5	172	179	23	0	0	374
2. C. F. Heaton New Burnside	3 lbs. Sulphate of ammonia	None	None	13	2	10	93	183	101	1	388
3. C. F. Heaton New Burnside	3 lbs. Sulphate of ammonia	Heavy in '30 & '31	None	13	3	83	183	97	12	3	378
4. M. J. McBride & Son, Villa Ridge	5 lbs. Nitrate of soda	Moderately heavy	None	14	1	8	152	222	40	0	422
5. C. F. Heaton New Burnside	None	Regular orchard mod. heavy	5"	14	10	1	13	73	104	18	209
6. M. J. McBride & Son, Villa Ridge	5 lbs. Nitrate of soda	Light cuts '30 & '31	5"	14	11	0	30	172	171	33	414

weeks or so before harvest. The last growth period is sometimes spoken of as the "final swell" and it is during this period that the effects of thinning are most noticeable.

Before passing to a more detailed consideration of the place of thinning in the orchard program, it might be well to note some general considerations regarding this practice. These have all been noticed previously by practical men and need only to be mentioned here in order to place them in the picture. To begin with, it is common knowledge that shoot growth

evened up. A reduction of the crop load can completely avoid breaking, except possibly with the older trees where winterkilling has injured the heartwood so that the scaffold limbs are brash. In this case bracing is necessary. An important after-effect of thinning is found in the greater hardness of both the tree and the fruit buds.

Thinning needs to be considered also in relation to nitrate applications, pruning, age of tree, type of growth, etc. The relationship between pruning,

(Continued on page 32)

AMERICAN FRUIT GROWER



*Sure...this lead
is Deadlier!*

Users of Chipman Hi-Test Lead Arsenate know for a fact that this lead is deadlier. They prefer it for codling moth control because:

IT'S PURER. Guaranteed to contain 98.15% di-ortho lead arsenate, recognized by authorities as the most desirable form.

IT'S Surer KILLING. Guaranteed to contain 32.50% killing ingredient (arsenic pentoxide), an 8% greater guarantee than ordinary leads guaranteeing only 30%.

IT'S SAFER TO FOLIAGE. Guaranteed to contain not more than 0.25% water soluble arsenic, which burns foliage. Other leads may contain up to 0.7%, practically three times as much.

Don't take chances with your fruit this year. Use Hi-Test Lead Arsenate and these equally effective Chipman Orchard Products: Dormant Oils, Sulphur Cream, Summer Oil, Dry Lime Sulphur, Nicotine Sulphate, Miscible Cresylic Acid.

Manufactured by Chipman Chemical Company, Inc., Bound Brook, N. J. Branch offices at Chicago, Ill., Houston, Tex., Tallulah, La., Palo Alto, Cal. and Yakima, Wash.

**CHIPMAN
HI-TEST
LEAD ARSENATE**

**Be a
McNess Man**

**No Time Like
Now to Get in—
Make up to \$75 a week**

It's no trick to make up to \$12 a day when you use your car as a McNess "Store on Wheels." Farmers are buying everything they can from McNess men. Attractive business-getting prizes, also money-saving deals to customers make selling McNess daily necessities a snap. This business is depression-proof.

We Supply Capital—Start Now!

There's no better work anywhere— pays well, permanent, need no experience to start and we supply capital to help you get started quick. You start making money first day. Write at once for McNess Dealer Book—tells all—no obligation. (92-A)

FURST & THOMAS, 456 Adams St., Freeport, Ill.

The GRAVELY
GARDEN TRACTOR &
POWER MOWER

Write for facts.
GRAVELY MFG. CO.
Box 603
Dunbar, W. Va.

Three machines in one...
Cultivates your garden...
Cuts the tallest weeds...
Mows your lawn... Ideal for Estate Owner.

These Tender Buds.....
YOUR NEXT
YEAR'S APPLES

Get PROTECTION this year without
sacrificing NEXT YEAR'S FRUIT!

use **KOLOFOG**
FUSED BENTONITE-SULPHUR

- 1 Because **KOLOFOG** is toxic but not caustic--it protects against diseases - - and allows the leaves to make big fruit and develop the buds - - YOUR NEXT YEAR'S APPLES.
- 2 Because **KOLOFOG** is colloidal, it mixes readily with water - - will not float on top or settle at the bottom of the spray tank - - thus assuring a uniform spray application over all fruit and foliage

Ask for Complete Information



NIAGARA SPRAYER & CHEMICAL CO., INC.

MIDDLEPORT, NEW YORK

SAN FRANCISCO, CALIFORNIA

JACKSONVILLE, FLORIDA

CANADIAN ASSOCIATE NIAGARA BRAND SPRAY CO., LTD. BURLINGTON, ONTARIO

NICHOLS

COPPER SULPHATE

99% pure, always uniform—the
standard for home mixed

BORDEAUX MIXTURE

Mono-Hydrated Copper Sulphate
35% Metallic Copper Content

NICHOLS COPPER CO.

A Unit of the Phelps-Dodge Corporation

40 WALL ST., N. Y. C. • 230 N. MICH. AVE., CHICAGO

EASTERN CHERRIES

(Continued from page 15)

fruit and side branches. Plenty of sunlight is also conducive to stronger growth and the forming of fruit-spurs on the lower branches where we wish to produce as large a portion of the crop as possible.

Annual or semi-annual pruning is recommended, for if pruning is delayed longer than this, the lower branches become shaded and weakened, and much of the bearing surface which will be produced at the top must later be cut away. All cuts are made to laterals growing out and downward, with plenty of bearing surface left throughout the tree. Sweets require much less pruning than sour, but are headed so that they may be picked with an 18-foot ladder. All sour are pruned so that they may be picked with an eight-foot stepladder.

Most of the harvesting is done by women and children, and if trees are regularly pruned, we have gone a long way to insure an ample supply of desirable help at harvest time. Thorough pruning also helps in maintaining a proper size and grade of fruit.

During the early years of the plantation it is advisable to intercrop with such row-crops as potatoes, cabbage, beans or tomatoes. The cultural practices employed in producing these interplanted crops are conducive to rapid development of the trees and their early maturity. After three or four years the orchard may be seeded down, preferably to a legume crop, and by the addition of manure or other mulching materials, maintained as a sod orchard.

Sweet cherries do their best under these conditions, and we find there is much less damage from cracking of fruit in the sod orchards than in the cultivated. Many of our best yielding Montmorency orchards have been down to sod for 15 years without having been broken up in that length of time. The grass which soon becomes a volunteer growth between the rows, is cut twice a season. Very little damage is caused by mice or rabbits to cherry trees. Spraying, harvesting, and all other operations are much easier when the orchard is maintained in sod. The result is more economical production.

Cherries, both sweet and sour, respond to liberal applications of nitrogen. Annual applications of from three to 10 pounds of nitrate of soda, or equivalent, depending upon the age and vigor of the tree, are applied about two weeks before blossom time. In order to maintain production, the tree should make a terminal growth of 10 to 14 inches each year. This growth develops the fruit spurs and forms healthy buds which are more

likely
temp
time

Th
whic
few
and
giver
all
Abou
phur
vious
contr
the a
which
plicat
made

Th
1300
550 a
apple
acres
cherry
Mont
lish
age f
planti
by 24
We
use a
pollin
Ou
expan
and M
are, l
sweet
one-h
and t
each
Ma
than
used
chard
ferti
My
terest
hire 2
chard
Our
three
spray
I be
organ
during
advert

Our
expan
and M
are, l
sweet
one-h
and t
each

Ma
than
used
chard
ferti

My
terest
hire 2
chard
Our
three
spray

I be
organ
during
advert

Tre

Per
trunk
spent
disea
shoul
woun
healin
Graft
plied
buddin

Pe

Two
Hubb
Coom
novelt
This f
and ap
only c
APRIL, 1937

plenty of
stronger
of fruit-
es where
a portion

uning is
g is de-
e lower
d weak-
ring sur-
at the top
cuts are
out and
ring sur-
Sweets
an sour,
may be
der. All
y may be
pladder.
done by
trees are
gone a
e supply
st time.
in main-
grade of

the plan-
rop with
cabbage,
ral prac-
these in-
to rapid
d their
or four
e seeded
ne crop,
nure or
aintained

st under
there is
cking of
an in the
st yield-
ave been
without
at length
soon be-
ween the

. Very
e or rab-
ng, har-
ions are
ard is
is more

sour, re-
of nitro-
of from
of soda,
n the age
applied
om time.

tion, the
growth
r. This
urs and
re more
APRIL, 1937

likely to withstand the frequent low temperatures just previous to blossom time.

The insects and fungous diseases which attack the cherry are relatively few and easily controlled. Leaf spot and maggot, with some attention given to brown rot, constitute about all the enemies of the Montmorency. About four sprays of lime and sulphur with arsenate of lead, three previous to picking and one after, will control these. The sweet cherry has the additional enemy of black aphid, which is easily controlled by the application of nicotine or other sprays made during the dormant period.

The present size of our farm is 1300 acres. The orchard acreage is 550 and is made up of 410 acres of apples, 100 acres sour cherries and 40 acres of sweet cherries. The sour cherry planting is about 90 per cent Montmorency and 10 per cent English Morello. These trees vary in age from two to 29 years, and the planting distance is 20 by 20 and 24 by 24 feet.

We hire bees for the orchard and use about one colony to the acre if pollination is a problem.

Our present plans call for a slight expansion of the sour cherry orchard and Montmorency will be used. We are, however, going to increase the sweet cherry planting, using about one-half Napoleon for the new stock and the remainder about one-quarter each of Windsor and Schmidt.

Manure obtained from our more than 100 head of stock on the farm is used for mulching in the cherry orchard. In addition to this we use fertilizers each year.

My father and brother are also interested in the farm operation. We hire 20 men by the year for the orchard, dairy and general farm work. Our mechanical equipment includes three trucks, seven tractors and 11 sprayers.

I believe the cherry industry needs organization for control of the pack during years of high production, and advertising for increased consumption.

Tree Wax Seals and Heals

Periodical examination of tree trunks may prove to be time well spent. Cracks, crevices, wounds and diseased tissue may be found which should be taken care of promptly. A wound dressing which is said to have healing qualities is now on the market. Graftwax-Tree Healtant can be applied at any time and is used also in budding and grafting.

Peachot Crop in Colorado

Two Colorado growers, Gene Hubbard of Orchard City and Fred Coombe of Cedaredge, harvested a novelty crop, peachots, last year. This fruit, a cross between the peach and apricot, is rare, as the trees bear only once every several years.

INSURE YOUR FRUIT CROP AGAINST

HAIL

One hail storm undoes the orchard work of months—destroys the brightest prospects for a profitable crop.


Why carry all the risk yourself? Why spend your time spraying, dusting, fertilizing, pruning, only to have it all wiped out by a single hail storm?

Your investment in money and time deserves protection in a reliable company writing Hail Insurance. Look at the list below, with risks distributed over the country, and let one of them relieve you of this heavy responsibility. Talk to your local agent now about Hail Insurance.

Aetna Insurance Company
World Fire & Marine Insurance Company
Piedmont Fire Insurance Company
Insurance Company of North America
Alliance Insurance Company
Philadelphia Fire & Marine Insurance Company
National Security Fire Insurance Company
Central Fire Insurance Company
Springfield Fire and Marine Insurance Company
Sentinel Fire Insurance Company
Michigan Fire & Marine Insurance Company
New England Fire Insurance Company

SEE LOCAL AGENT IN YOUR TOWN
OR ADDRESS THE COMPANY

HAIL DEPARTMENT—209 W. Jackson Blvd., Chicago, Ill.



APPLES RUINED BY HAIL



THESE APPLES A TOTAL LOSS

RUNNING WATER FOR EVERY FARM

COMPLETE SYSTEMS
from 200 gallons to
5,000 gallons per
hour at new
low price

FREE BOOK tells how you
can have run-
ning water all year 'round for
only a few cents a day. Just
mail the coupon.



Fairbanks, Morse & Co.
900 S. Wabash Ave., Dept. 4791
Chicago, Illinois
Please send me your illustrated free book on water systems.

My source of water supply is:
Lake..... Shallow Well..... Stream
Spring..... Deep Well..... Cistern
Have you electricity? Yes..... No

Name.....
Address.....
Town.....
R.F.D.....
State.....

Whether or not you have electricity, whether you draw from deep or shallow well, lake, or stream, you can have running water at remarkably low cost. Complete systems, from 200 gallons to 5,000 gallons per hour at new low price.

FAIRBANKS-MORSE HOME WATER SYSTEMS

Engine or motor driven for every farm or home use





AUDITORIUM

HOTEL RATES

Now from \$1.50

Just think—you can now enjoy the comfort and luxury of the Auditorium Hotel for as little as \$1.50 a day for a room without private bath or from \$2.50 a day with private bath. A splendid location—every convenience—in an atmosphere of luxury at low cost.

George H. Mink, Manager

MICHIGAN AVE. AT CONGRESS

CHICAGO

*When worms +
scab come round to call,
Hamilton Guns will
kill them all
Will Hamilton*

Spray Guns with Controlled Streamline

W. L. HAMILTON & CO.
BANGOR - MICHIGAN

Six Models - A gun for every purpose

WINE MAKERS MANUAL


In this book Peyton Boswell reveals every secret he employs at U. S. Winery No. 59 which will benefit the home wine maker or the small winery. Whoever follows the methods described cannot fail to make the best possible wine from the variety of grapes he uses. Written in easy, understandable style. Illustrated.

Sent postpaid on receipt of \$1.50

AMERICAN FRUIT GROWER

1370 Ontario Street Cleveland, Ohio

Stoves AT FACTORY PRICES!



SALE CATALOG—FREE

Nearly 200 Styles and Sizes of Heaters, Ranges, Furnaces at Factory Prices. Easy Terms—as little as 18c a day—Year to Pay. More Bargains than in 20 Big Stores. New styles, new features, new colors. 30 days free trial—360 days approval test—24-hour shipments. The Kalamazoo Stove Co., Manufacturers, 495 Rochester Avenue, Kalamazoo, Michigan.

Over 1,100,000
Satisfied Users
37 Years in Business
Write for FREE Catalog

**"A Kalamazoo
Direct to You"**



Two limbs with about the same "leaf surface" which shows the reduction in size from "clustering" but a greater yield.

PEACH THINNING

(Continued from page 29)

nitrating, and thinning has been given some attention in the Illinois experiments. One page from our notebook has been included in Table I, which illustrates what can or cannot be done to the crop through these three variables.

Since the objective in growing peaches is to obtain both size and yield, the data summarized will be particularly interesting from that angle. Of the salient points which may be noted in the table, the unthinned trees in the Heaton orchard (Experiment 1) are of special interest. In this instance the unthinned tree matured an average of 374 pounds of fruit per tree but a large proportion of the crop was below one and three-fourths inches in size. These particular trees matured an average of 3,200 fruits each and with this much of a load it was impossible to size up the crop even though the yield was not excessively high. By way of contrast, the trees in Experiment 6 may be noted. Here the yield was higher but the number per tree was limited. A large proportion of the yield fell in the larger commercial sizes.

The success of sizing up fruit through nitrate applications when pruning and thinning are omitted is illustrated in Experiment 2. In this instance the upper limit of a crop load was reached in the absence of thinning and as a result much of the crop, particularly that in the clustered part of the tree, fell into the smaller sizes. By way of contrast, in Experiment 3, heavy pruning and nitrating were tried without thinning, with a result even more disastrous than in the above. Likewise in Experiment 4 with fairly heavy nitrate applications and

with moderately heavy pruning, thinning was omitted. The yield was high, but as in the other unthinned trees, the size was small. In Experiment 5 the crop was limited by both thinning and pruning but nitrogen was omitted. In this instance the fruit set lighter and tended to drop to a greater extent, thus limiting the load to nearly half that of some of the other treatments. In these trees there was a type of growth, generally spoken of as a high carbohydrate growth, which produced an especially fine color in the fruit although the yield was cut. The fruit ripened evenly and was all harvested at one picking.

The general trend of the results in this table will show definitely how these three variables can be used in controlling size. Judgment will be required to direct the type of thinning required under different growth conditions but with experience these variables can all be brought into balance and the stage set each year for a fair yield even if growth conditions happen to be unfavorable.

One more thing now needs to be considered and that is the "final swell." The thinning problem as related to the other treatments, comes to a focus at this time. Experienced growers know how difficult that last week is with a peach crop. Always the possibility of prolonged rainy periods with brown rot and delayed picking have to be considered, to say nothing of the price level. In other words, when shall we pick? If the picking is done too soon, both quality and yield are cut. If done late the quality increases but the shipping radius is shortened. Studies at Illinois station show that a peach enlarges when the

moisture supply is adequate up until two or three days, after it starts to become soft. Obviously, then, overloading becomes more acute during the final three or four days than at any other time. This point has been carefully considered at a number of stations and viewed from the marketing angle it seems that peach growers will have to give more thought to the degree of ripening before picking is started.

With a 25-foot planting distance, an average yield of mature trees could easily be placed at 300 to 500 bushels per acre. During the last week if growth conditions are adequate and the crop load not too heavy, peaches enlarge in the neighborhood of one-sixteenth of an inch or more in diameter per day. In other words, expressed in the terms of bushels, the yield increases five to fifteen bushels each day. This fact can be taken full advantage of by careful picking or "topping." Picking too soon may be serious from the standpoint of the bearing of quality upon repeat orders. On the other hand, the time element becomes an important factor where large acreages are involved. When all of these variables are taken into consideration, it requires good management to control the situation. Other things being to our choosing, picking can be gaged by the shipping distance and the market requirement as to ripeness. At best, however, the time is so short, and the time limits so great, that everything has to "click" at harvest time.

There is another aspect to the size problems in peaches which needs to be taken into consideration. My associate in these experiments—R. L. McMunn—has determined that there is about twice as much surface to be peeled in a bushel of one and three-fourths-inch peaches as compared to a bushel of three-inch peaches. And as strange as it may seem, there is more flesh in 50 pounds of peaches of the larger size than in the smaller, partly because there are fewer seeds.

In its final analysis, it will be seen that there are a good many angles to the thinning problem as it is related to the tree condition and the crop load. It would seem that under eastern conditions it would be only fair to set the stage each year for average growth conditions. If the season turns out to be especially dry, the commercial sizes can be reached with a fair proportion of the crop. Thinning is thus seen to be more necessary during those years when a large crop is in prospect, so that the general level of the pack can be raised accordingly.

APRIL, 1937

MAGNETIC SPRAY

the Sulphur that gives
MAXIMUM CONTROL

Every year more growers are learning of the economy of using Stauffer Sulphurs. They go farther, and give longer and better protection against scab and other fungous growths. Use Stauffer Sulphurs this season and see why more growers prefer it to all others. There's a grade for every purpose and for every type of equipment.

"MAGNETIC SPRAY" WETTABLE SULPHUR. The Sulphur in the spray does the work—the higher the sulphur content the more effective the spray. Insist on Magnetic Spray—98.5% Pure.

"MAGNETIC" CATALYTIC SULPHUR. A safener that stops the damaging reaction that takes place in lead arsenate-lime sulphur sprays. It also should be used in straight lime sulphur solutions.

"ELECTRIC" SUPER-ADHESIVE DUSTING SULPHUR—99.8% Pure. When you use a dust—use "Electric"—and be sure.

OTHER MAGNETIC PRODUCTS

Rotenone-Sulphur Dust
Pyrethrum-Sulphur Dust
"Alorco" Cryolite
Cryolite-Sulphur Dust
Cryolite-Sulphur Spray

STAUFFER CHEMICAL COMPANY
420 Lexington Ave., New York City
Chicago, Ill. Freeport, Texas



Stauffer Chemical Co., 420 Lex. Ave., N.Y.C.

Please send the following literature

"Magnetic Spray" ☐
Wettable Sulphur ☐
"Electric" Dusting ☐
Sulphur ☐

"Magnetic" Catalytic ☐
Sulphur ☐

I grow _____

Name _____

Address _____



TEMPERED RUBBER

GIVES THE "U.S." ROYAL BOOT ONE THIRD LONGER WEAR THAN ORDINARY BOOTS. DISTINCTIVE TIRE TREAD SOLE. PIGSKIN FINISH. AND, LIKE OTHER "U.S." BOOTS, EVERY BOOT IS LEAK-TESTED BEFORE IT LEAVES THE FACTORY.




UNITED STATES RUBBER PRODUCTS, INC.
1790 BROADWAY NEW YORK

ASK TO SEE THE
NEW ROYAL RAINCOAT
WITH PIGSKIN FINISH

United States Rubber Company

AMERICAN FRUIT GROWER

PAGE 33



"SELF-SERVER"

Apple and Pear Cutter

The perfect fruit sampler for growers. Let your customers and buyers sample your product. The perfect purchase or gift for fruit consumers.

Beautifully and Strongly Made
Quadruple Silver Plated
Will Last a Lifetime

\$2.25 postpaid

AMERICAN FRUIT GROWER

1370 Ontario St., Cleveland, O.

CHARACTER

Here, one instinctively feels those perfections of hospitality and service that have made The Bellevue famous . . . Rates Begin at \$3.50

BELLEVUE STRATFORD

PHILADELPHIA

One of the few famous hotels in America

CLAUDE H. BENNETT, Gen. Mgr.

ENJOY BEAUTIFUL FALSE TEETH



SEND NO MONEY
UNITED STATES DENTAL COMPANY
Dept. 405 1557 Milwaukee Avenue Chicago

CROTALARIA SPECTABILIS

World's Finest Summer Legume

Build up soil. Prevent erosion.

SIMPSON NURSERY CO., Monticello, Florida

We sold the seed to Mr. Lutz whose article appeared in December issue of AMERICAN FRUIT GROWER.



STRAWBERRIES

Allen's 1937 Berry-Book Describes Best Methods, Plants, Varieties:

Fairfax, Dorsett, Catahill, etc.

COPY FREE WRITE TODAY

THE W. F. ALLEN CO.

284 Evergreen Ave., Salisbury, Md.

NEW KINKADE GARDEN TRACTOR

and Power Lawnmower

A Practical, Power Plow and Cultivator for Gardeners, Suburbanites, Florists, Truckers, Nurserymen, Fruit Growers

Low Prices - Easy Terms

American Farm Machine Co.

1105 33rd Av. SE, Minneapolis, Minn.

PAGE 34

PROFITABLE PEACH PRODUCTION FACTORS

(Continued from page 10)

We think that the poorer growth made by the trees in lespedeza and sweet clover is due to the fact that they suffered more in periods of moisture deficiency than trees in cultivation alone or with summer cover crops. From moisture samples taken, we observed that the amount of moisture in the soils under the sweet clover and lespedeza was reduced to a critical point, that is, to the wilting percentage of moisture for that soil, for a considerable period of time in each of those years. In 1936 we had little summer rainfall; the moisture in the first two feet of soil in the lespedeza plot was reduced both by the trees and the cover crop to the wilting percentage for a four weeks' period in late June and July, and again in August for a three weeks' period from August 1 to August 21. This last period was the time of final fruit swelling.

The growing cover crops require plant nutrients for their growth and in this way compete with the tree. In 1936, we studied the nitrate content of the soil of plots at weekly intervals. Nitrates were low under lespedeza and cultivation, but high under sweet clover and soybeans. The growth of the trees, therefore, appears to have been affected more by periods of critical moisture shortage rather than by nitrate deficiency.

We have been attempting to determine the best way to handle the sweet clover-lespedeza to reduce to a minimum this competition with the trees for moisture. It may mean that alternating middles of lespedeza or sweet clover with a summer cover such as soybeans, cowpeas, or crotalaria will be satisfactory. Whatever the cover used, it should be clipped or disked up during long dry spells to conserve the moisture for the trees.

There are certain definite advantages in the use of cover crops in a peach orchard. Where soils are not of sufficient depth to permit the use of summer cover crops, certainly a winter cover crop should be used, not only from the standpoint of its benefit in preventing washing, but also from the standpoint of its nutritional value to the trees when disked under each spring. The cover crop serves a very important function in utilizing mineral elements, particularly nitrates, and preventing their leaching from the soil in winter. These are made available to the tree along with other important mineral constituents when the cover crop decays in the soil.

While the practice of growing summer and winter cover crops may be used to advantage during the early life of the orchard, the single winter cover crop may be the only one that

can be used after the trees have reached bearing age. Since in many orchards the trees are planted 20 by 20 feet apart, the root systems come to occupy all of the area between the trees when the trees are about five to six years of age. Cover crops, especially deep-rooted ones like sweet clover, may compete with the trees when this stage of growth is reached.

Many growers report that they are unable to obtain even a winter cover crop in their old peach orchards after the trees have begun to crowd. A satisfactory cover will not be obtained unless fertilizer is used at the time the cover crop is seeded. I have seen the production of very satisfactory cover crops in old orchards by this method.

Another very important use of the cover crop in addition to its nutritional value is to facilitate water penetration and prevent run-off. In our experimental orchard which is planted on a rather steep slope, very little washing has been observed from spring rains following disking in of a good rye and vetch winter cover crop. Likewise, on this same soil, little washing has been observed from late summer rains after good summer cover crops have been disked under. By a good cover crop, of course, is meant a good growth of plants over the entire area.

Other important factors in the profitable production of peaches are the size and quality of fruit. It is in years of heavy production especially that these factors are important in bringing an increased price to the producer. Our most important commercial variety of peach which we are growing today is not one of the highest quality even when grown under favorable conditions. On overloaded trees with poor leaf area, where sufficient thinning has not been done, the fruits may be small and of inferior quality. This kind of fruit does not sell readily and in years of heavy production fruits of poor size and quality do not appeal to the consumer, and their accumulation on the market tends to aggravate the price depression created by a large crop. In Virginia the Elberta variety has proved very satisfactory as a commercial variety. It fits well into the seasonal distribution of peaches from other producing centers. Here, as in other states, however, this variety is not of the highest quality when produced on trees with a poor leaf area and on soil inadequately supplied with moisture.

In experimental studies made with the Elberta variety, we have found that fruits grown with a small leaf area—about 10 to 15 leaves per fruit—are small, of rather poor quality,

low in s
With a m
to 40 lea
eating qua
the necess
just the f
the set is
Pruning
load as w
of bearing
adjunct to
and qualiti
maintainin
been ment
trees is al
in stimulu
growth.
near the
usually be
these sho
so that the
exposed t
mum lea
duced are
color.
An ad
available
portance
color and
fruit exh
periods as
blossomin
berta vari
about 60
increases
suture .di
rather slo
which is
hardening
days. Fin
is noted v
before fin
This la
knows as
one of ma
the fruit
When soi
this final
fruits on
thinned, th
fruit size
condition
of Virgin
for the m
small size
cover cro
during thi
to be cons
soils.
In addi
may affec
portance
varieties
As a resu
has been
years, we
results in
ties of hi
have been
others are
show dis
who are
ties that
Peach gr
be intere
which m
APRIL, 1937

AMERICAN FRUIT GROWER

APRIL, 1937

low in sugar and somewhat bitter. With a more adequate leaf area—30 to 40 leaves per fruit—the size and eating quality are far superior; hence the necessity for hand thinning to adjust the fruit-leaf ratio in years when the set is heavy.

Pruning to regulate the size of the load as well as to increase the vigor of bearing trees is also an important adjunct to thinning in securing size and quality. The value of nitrogen for maintaining tree vigor has already been mentioned. Pruning of bearing trees is also important and necessary in stimulating a vigorous shoot growth. Vigorous shoots produced near the ends of cut-back branches usually bear large leaves, and when these shoots are thinned sufficiently so that the leaf surface is well enough exposed to sunlight to permit maximum leaf function, the fruits produced are usually of good size and color.

An adequate supply of moisture available to the tree is of equal importance to leaf area in obtaining color and size of fruit. The peach fruit exhibits three definite growth periods as it increases in size from blossoming to maturity. For the Elberta variety, the first period covers about 60 days while the young fruit increases principally in axial and suture diameters. Then follows a rather slow period of development which is roughly the period of pit hardening and which lasts about 40 days. Finally a third period of growth is noted which starts about 30 days before final harvest.

This last period which the grower knows as the period of final swell is one of marked increase in volume, and the fruit fills out in cheek diameter. When soil moisture is lacking during this final period, even though the fruits on the tree have been well thinned, the unfavorable effect on final fruit size will be marked. It was this condition that existed in many parts of Virginia this year and accounted for the large quantity of fruit of small size and color. The effect of cover crops on the moisture supply during this period of final swell is one to be considered, especially on shallow soils.

In addition to growth factors which may affect size and quality, the importance of developing higher quality varieties of peach must be emphasized. As a result of the breeding work that has been in progress for the past 25 years, we are just beginning to see results in the production of new varieties of higher quality. Some of these have been fairly widely tested while others are on preliminary trial. Many show distinct promise for growers who are interested in planting varieties that mature earlier than Elberta. Peach growers of any region should be interested in these new varieties which may compete on the market

with the varieties they are now growing. It is a safe prediction that commercial peach varieties of the future must be of higher quality if this fruit is to hold its place on the market in competition with other fruits.

POLLINATION

(Continued from page 16)

len was carried 200 miles to the largest planting of Eumelan vines in New York State and there mixed with Lycopodium powder, also a pollen product. Several hundred vines of Eumelan were then dusted with the mixture, and a like number untreated were left for a comparison. The time consumed in the operation was not over an hour.

Largely because this variety had borne very light crops the past few years, and as a result plant food reserves had been stored in excess of the quantity normally needed, the clusters were more compact the past season than usual. It has already been noted that well-nourished vines, even though they be self-sterile or imperfectly-fertile varieties, form better clusters when there is a sufficiency of plant food.

In spite of the fact that the clusters were above normal in compactness, these vines that were cross-pollinated with the Aramon X Rupestris Ganzin pollen, using the hand duster for dispersal of the pollen, yielded 50 per cent more fruit than the untreated. The increase was due to the fact that more berries on the cluster were fertilized on the dusted than on the untreated vines which were dependent on chance cross-pollination and fertilization.

In view of these facts, it seems entirely practical from a commercial standpoint to cross-pollinate artificially certain varieties of grapes that are greatly in demand for specific purposes, thereby improving the appearance of the clusters for dessert purposes and at the same time greatly increasing the yields.

APRICOT CULTURE

(Continued from page 27)

ern market are usually packed in standard four-basket crates, weighing packed approximately 26 pounds; 1,000 crates generally constitute a carload. Of more recent years an increasing tonnage has been shipped either packed or loose in small lugs of about the same capacity as the crate.

Apricots are sold to the cannery at a given price per ton, the price being determined to a certain extent by size which is specified as so many fruits per pound.

AMERICAN FRUIT GROWER

FLAME-GUN DESTROYS WEEDS

Does Work of 4 Men

Hauck Kerosene Flame-Gun—2000°F. controlled heat—quickly...easily...economically destroys weeds (seeds and all), brush, other objectionable growths. Keeps roadways, fences, irrigation ditches, orchards, etc., clean. Hundred and one uses. Inexpensive—Safe—Easy to use. Pays for itself in time and money saved. 10 day Free Trial. Write for Free literature and special introductory price.



HAUCK MFG. CO.

161 TENTH STREET
BROOKLYN, N.Y.

FREE VALUABLE FOLDER—WRITE TODAY!

BAND YOUR APPLE TREES

KILL CODLING MOTH

Our experience has taught us how to chemically treat **TREE BANDS** that are guaranteed to kill the worms.

BANDING saves one or more spray applications—does away with stings—and kills thousands of worms that would live over winter. Costs less than 2 cents per tree.

Write for Circulars and Prices

EDWIN H. HOUSE

Saugatuck, Mich.

Remember we pay the freight.

Jim Brown's Factory Prices on FENCING

Will Save You Big Money

Amazing New Process, using COPPER Steel Wire makes my fence last much longer than ordinary fence. Don't buy fencing till you get my new Factory Prices.

Jim Brown Pays the Freight

Write factory nearest you for catalog of bargains in farm and home needs—Fencing, Gates, Steel Posts, Barb Wire, Baby Chicks, Poultry Supplies, Stoves, Harness, etc.—Jim Brown, 65

THE BROWN FENCE & WIRE COMPANY
Dept. 165 Cleveland, Ohio; Memphis, Tenn.

MAN WITH CAR WANTED



To conduct world-renowned home service business—countrywide. Nothing new. Now over 8,000 Rawleigh Dealers. Many do \$3,000-\$5,000 annual business in necessities for home-larm. Stocks supplied on credit. Write for information how to start in business on our capital. Dept. D-70-AFG.

W.T. Rawleigh Co., Freeport, Ill.

*Includes nationally recognized D. & D. for orchard spraying—thousands of gallons sold yearly.



Kill Garden Weeds QUICK - EASY - with a Barker Weeder, Mulcher & Cultivator.

"Best Weed Killer Ever Used." Makes perfect soil mulch at same time. Only weeder-mulcher with Filler Drum. 50% better. Easiest to use; lowest prices. 8 sizes. Write

BARKER MFG. CO., Box 2, David City, Nebr.



NEW SUPER-QUALITY KITSELMAN FENCE

Heavier, tougher pure zinc galvanizing. Stronger, tougher Copper-Bearing Steel. Same low Factory Prices. WE PAY FREIGHT. 160 styles and heights Farm, Poultry and Lawn Fence; Steel Posts, Gates, Barb Wire, Paints, Roofing. Write today.

KITSELMAN BROTHERS
Box 206 Muncie, Indiana

GRAFTWAX-TREE HEALANT is waterproof—adhesive. Used cold right from can. Gives nearly 100% success in grafting and budding; arrests and cures Tree and Plant Diseases (including blights). Excels as Wound Dressing for GRAFTWAX SEALS AND HEALS. ¼ lb., 50¢; 2 lbs., \$1; 6 lbs., \$3; 12 lbs., \$5.—Postpaid. Send for free sample and Testimonials.

CLARION DEVELOPMENT CO., INC.

Dept. A, Clarion, Pa.

PAGE 35

OPPORTUNITY ADS

Only 15c a Word—CASH WITH ORDER
Count each initial and whole number as one word
ADDRESS: AMERICAN FRUIT GROWER,
1370 Ontario Street, Cleveland, Ohio

BABY CHICKS

AS WORLD'S LARGEST CHICK PRODUCERS. WE can save you money. Big Catalog Free. COLONIAL POULTRY FARMS Pleasant Hill, Missouri.

BEES

PACKAGE BEES WITH PURE BRED ITALIAN Queens 2 lb. package \$2.45, 3 lb. package \$3.15. Place orders early for service. RAPIDES APIARIES, Winnfield, Louisiana.

FULL 8 FRAME COLONIES, IN STANDARD HIVE bodies \$6.00. March, April shipments. MILLERS APIARIES, Three Rivers, Texas.

BEES FOR CROSS POLLINIZING FRUIT BLOOM OR raising honey. Write for prices. D. C. JACKSON, Funston, Georgia.

BERRY BOXES

USE MODERN GREEN AND WHITE ROLLER BERRY Boxes for Fruit and Vegetables. Write now. Liberal Discounts. ROLLER BOX COMPANY, Dept. E, Benton Harbor, Michigan.

BERRY PLANTS

BOYSENBERRY, ALSO FIRST RELEASE OF NEW FRANKLIN D. AND NECTARBERRY PLANTS, which yielded for us last year twice the fruit of Boysenberry. Thornless Loganberry, Lloyd George Raspberry and others. BENEDICT RANCHO, 1937 Deana Road, El Monte, California.

MECHANICAL STRAWBERRY TRANSPLANTER. Write in at once for literature and information about this new Mechanical Transplanter proven with great advantage in many sections for years. HOLLAND CELERY PLANTER COMPANY, Holland, Michigan.

RANCHO VERDAD BOYSENBERRY. WORLD'S largest vineberry. Taste resembles raspberry. Hardy, true to name plants prepaid 10—\$2.00, 100—\$12.00. RANCHO VERDAD, Mentone, California-Ardmore, Oklahoma.

STRAWBERRY PLANTS—CERTIFIED: AROMA, Blakenore and Missionary 500, \$1.50; 1000, \$3.00; 5000, \$13.50. N. T. BASHAM, Mountainburg, Arkansas.

CIDER MILLS

MAKE A PROFIT ON YOUR CULL APPLES BY PUTTING them into sweet cider. Presses, graters, pumps, filters, supplies. Booklet F Free how to keep cider sweet, make vinegar quickly, and money in roadside mills. PALMER BROS., Cos Cob, Connecticut.

CROTALARIA SEED

CROTALARIA SPECTABILIS SEED: HEAVIEST LEGUME cover crop. Reserve now. GRAND ISLAND NURSERIES, Eustis, Florida.

DAIRY GOATS

DAIRY GOAT JOURNAL, DEPT. 703, FAIRBURY, Nebr. Monthly Magazine, 50c yearly. Introductory 3 months 10c.

HOSIERY

BEAUTIFUL SILK HOSIERY 5 PAIRS \$1.00. FULL-fashioned 3 Pairs \$1.00 (Trial 39c pr.). DIRECTCO, AF221W Broad, Savannah, Georgia.

NURSERY STOCK

FRUIT TREES AND BERRY PLANTS FOR SPRING Planting. 200,000 Peach Trees, 150,000 Apple Trees, one and two year old. Pear, Plum, Cherry, Nut Trees of all kinds. Millions of Strawberry, Raspberry, Blackberry, Asparagus Plants, and Grape Vines. Best new and old varieties. Evergreens, Shade Trees, Shrubbery and Roses. We offer one of the largest and most complete lines of Nursery Stock in the East, sold direct to the planters at comparatively low prices. Send today for our free catalog. BOUNTIFUL RIDGE NURSERIES, Box H, Princess Anne, Maryland.

HARDY ENGLISH WALNUTS—CRATH'S CARPATHIAN strain. New. Parent stock hardy at forty degrees below zero. Descriptive pamphlet free. S. H. GRAHAM, 42 W. Bostwick Road, Ithaca, New York.

NEWBURGH—THE BIGGEST RED RASPBERRY. Mosaic-free. Special price, No. 1, \$17.50 thousand; No. 2, \$9.00 thousand. McDonald Red Rhubarb, outstanding, \$17.50 hundred. FARM OF PASCHKE, Northeast, Pennsylvania.

HARDY APPLE AND PEACH TREES, SEEDLINGS and Root Grafts. Write us. JONES NURSERY COMPANY, Woodlawn, Virginia.

ORCHARD MANAGER WANTED

GOOD, HONEST, CAPABLE, for 80-acre orchard. Must know orcharding. ANNETTE B. EHLEB, Hennessey, Oklahoma.

PAGE 36

PATENTS

National Trade Mark Company
Munsey Building
Washington, D. C.
Trade Mark Specialists

PATENTS. LOW COST. BOOK AND ADVISE FREE. L. F. RANDOLPH, Dept. 563-A, Washington, D. C.

PHOTO FINISHING

FILMS DEVELOPED AND PRINTED, INCLUDING two beautiful olive tone enlargements Free. 25c Coin. UNITED PHOTO SERVICE, La Crosse, Wisconsin.

FILMS DEVELOPED ANY SIZE. 25c COIN, INCLUDING two enlargements. CENTURY PHOTO SERVICE, Box 829, La Crosse, Wisconsin.

PLANTS FOR SALE

FROSTPROOF CABBAGE, EACH BUNCH FIFTY, mossed, labeled variety name, Jersey Wakefield, Charleston Wakefield, Succession, Copenhagen, Early and Late Dutch, postpaid: 200, 65c; 300, 75c; 500, \$1.00; 1,000, \$1.75. Onion: Crystal Wax, Yellow Bermuda, Sweet Spanish, Prizetaker, Prepaid: 500, 60c; 1,000, \$1.00; 6,000, \$3.50. Tomato: Large, well rooted, open field grown, mossed, labeled with variety name, Livingston Globe, Marglobe, Stone, Baltimore, June Pink, McGee, Earliana, Gulf State Market, Early Detroit, Postpaid: 100, 50c; 200, 75c; 300, \$1.00; 500, \$1.50; 1,000, \$2.25. Pepper mossed and labeled, Chinese Giant, Bull Nose, Ruby King, Red Cayenne, Postpaid: 100, 65c; 200, \$1.00; 500, \$1.75; 1,000, \$2.50. Full count, prompt shipment, safe arrival, satisfaction guaranteed. UNION PLANT COMPANY, Texarkana, Arkansas.

TOMATO PLANTS. GARDENERS GROW RIPE TOMATOES two weeks earlier with our hardy, field-grown tomato plants. Leading varieties grown from best Certified Seed. \$1.50 per thousand. Largest Individual Grower. 1500 Acres in Vegetable Plants. Write for list. CARLISLE PLANT FARMS, Valdosta, Georgia.

PIEDMONT PLANTS—WAKEFIELD, COPENHAGEN, Golden Acre, Flat Dutch Cabbage, Bermuda, Valencia, Prizetaker Onion, Postpaid: 200, 50c; 500, \$1.00; 1000, \$1.75. Collect: \$1.00 per 1000. Earliana, Bonny Best, Pritchard, Marglobe, Baltimore Tomato Plants, ready April 15th. Postpaid: 300, 80c; 500, \$1.50; 1000, \$2.50. Collect: \$1.75 per 1000. Write for catalogue of many other vegetables and flower plants. Satisfaction Guaranteed. PIEDMONT PLANT COMPANY, Albany, Georgia.

GROW VEGETABLES THREE WEEKS EARLIER WITH Carlisle's hardy, Field-Grown plants. Cabbage plants, \$1.00. Onion, 60c; Tomato, \$1.50, thousand. Satisfaction guaranteed. Catalogue free. CARLISLE PLANT FARMS, Valdosta, Georgia.

C.O.D. SEND NO MONEY. FROSTPROOF CABBAGE and Onion Plants. All varieties. 500, 50c; 1,000, 90c. Prompt Shipment. GEORGIA PLANT COMPANY, Albany, Georgia.

HARDY, FIELD-GROWN PLANTS—BROCCOLI, Cauliflower, Cabbage, Onion, Eggplant, Tomato, Pepper, Sweet Potato Plants, Moss Packed. Shipped anywhere. Safe arrival guaranteed. Largest Individual Grower. 1500 Acres Vegetable Plants. Descriptive Folder. CARLISLE PLANT FARMS, Valdosta, Georgia.

POROUS IRRIGATION HOSE

IRRIGATION PAYS—WRITE ABOUT POROUS HOSE irrigating. B. & B. IRRIGATING SYSTEM, Port Clinton, Ohio.

PRINTING

POSTPAID! LETTERHEADS, ENVELOPES, 500 EACH Printed \$3.50. 250 Business Cards \$1.00. SEE-JAY, 2459 Station St., Chicago, Illinois.

PRUNING EQUIPMENT

GUARANTEE SAVE 50% ON TREE TRIMMING WITH simple, light invention. No climbing. Does job successfully in one-half time. CLIPPER SAWS, 317 N. Noble Street, Indianapolis, Indiana.

SERVICES

TREE GRAFTING AND BUDDING. HOWARD H. MARTIN, Princeton, Illinois.

SPRAYER VALVES

JOBBER AND GROWERS WRITE FOR PRICES AND catalog. Complete line of Sprayer Valves. CENTRAL BRASS MANUFACTURING COMPANY, 2950 E. 55, Cleveland, Ohio.

TREE BANDS

BETA-NAPHTHOL TREATED TREE BANDS "SURE KILL" the codling moth worms. Write for prices and free literature. M. A. KOELLER, Barry, Illinois.

AMERICAN FRUIT GROWER

NUT VARIETIES

THE results of a survey conducted by D. C. Snyder of Iowa, chairman of the survey committee of the Northern Nut Growers' Association and presented at the meeting of the Northern Nut Growers' Association at Geneva, N. Y., in September, 1936, revealed considerable information concerning many of the nut varieties discovered in the earlier prize nut contests of the association. These reports come from a wide range of conditions and with the older sorts cover several years' observations. The prospective planter of nuts will find this report very instructive.

Thomas, one of the oldest of the black walnut varieties, was mentioned favorably by nearly all of the 64 growers who reported. Its most serious defect is susceptibility to the canker disease. The Ohio black walnut was generally unsatisfactory, being rather unproductive, lacking in quality and with too thick a husk. Stabler also received many unfavorable reports, being lacking in hardiness, vigor and quality.

Stambaugh was reported favorably by many and was considered by Mr. Snyder as being a valuable variety for some sections, the best reports coming from Illinois and Oklahoma. Ten Eyck, another of the older varieties, also received favorable mention from several growers. Many other black walnut varieties were listed in Mr. Snyder's paper, but information concerning them was too meager to warrant any conclusions as to the merits of the varieties.

The Winkler native hazel was generally satisfactory within a radius of 200 miles of its place of origin in southeastern Iowa. Rush, another native hazel, received many favorable reports, although there were some reports of its lacking hardiness. For a native hazel it is about as good as any of its class now available.

Many hickories were mentioned, but the information concerning any one variety was very meager. The Taylor, Kirtland and Kentucky hickories were considered to be three of the best by Dr. W. C. Deming whose knowledge of hickory varieties is very extensive. Stratford was considered promising because of its early bearing habit and adaptability to bitternut stocks. Rockville also received favorable mention by several. Fairbanks was hardy, vigorous and productive, but only medium in quality. Swain was also considered good.

Certainly one who reads this report receives the impression that the need is great for more extensive variety tests. The need is equally great for better varieties.—G. L. SLATE, Sec'y, Northern Nut Growers' Assn., Geneva, N. Y.

EDITORIAL

(Continued from page 7)

the hazards are somewhat reduced over former times. A block of prunes, sweet cherries where they will thrive, the more disease-resistant raspberries, and the superior new strawberries should attract many customers to the roadside market and embellish the load which goes to the city market. All this means a better American horticulture and a more profitable industry.

Again, there is the desirability in some instances of diversifying with poultry, dairying, or some other line than fruit. Poultry fits into the picture quite well, provided the market is not already oversupplied. It requires an entirely different equipment and knowledge but in nearly every section one finds these two lines successfully carried. But as a rule the general farmer has not been a very successful orchardist and for the large scale farmer to have a few acres of orchard is quite the reverse of the thought here expressed. The farm orchardist has rapidly passed out and the intensive fruit specialist has taken his place. But with this new situation a certain amount of diversification has its place.

APRIL 1937

NEW


By HANDY ANDY

Comes April and the busy season, but the really busy fruit grower is never too busy to stop and pick up a new idea. This is the place to exchange some of these ideas and to let each other know about new products and ideas for our mutual benefit. You probably have some ideas that are new and helpful so just drop me a line about them. The address is Handy Andy, American Fruit Grower, 1370 Ontario St., Cleveland, Ohio.

HIGH SPEED PUMPS •

Orchard irrigation calls for plenty of water. Frank H. Wissler, Virginia fruit grower, says that he has used as high as 2,000 gallons of water per tree every two weeks during the growing season. To supply the large amounts of water for irrigation it is necessary to have a pump that will give the proper delivery. I know of one fruit grower who obtains his water from a river that is more than 75 feet below the level of his orchard.

New, high-speed centrifugal pumps are now on the market that will deliver from 10 to 6,000 gallons per min-



ute at a head of 10 to 280 feet depending on the size of the unit. The new pump unit usually includes a coupled electric motor of enough power to suit the size of the pump. Equipment like this provides water for every part of the orchard whether large amounts of water are needed for irrigation or for filling storage tanks for spraying water. The centrifugal pump is particularly adapted to pumping water from rivers and reservoirs. Other types are available for wells.

It seems water is becoming more and more important on the fruit farm. The latest use for water in large quantities is for washing the fruit to get rid of residues. The U.S.D.A. recommends that each bushel of fruit put through the acid bath of the

- High Speed Pumps
- Air Conditioned Carton
- Carboy Cart and Tilter

washer be rinsed with three or four gallons of clean water. That means a lot of water even for the medium-sized orchard.

AIR CONDITIONED CARTON •

Here's a new air conditioned fruit shipping box that cradles each fruit in a separate compartment. The box




itself is made of sturdy corrugated fibre board and is constructed with air vents so that the air can circulate through the box. This type of ventilation allows for rapid escape of field heat resulting in less damage from breakdown and other high-temperature troubles. As you can see from the illustration, the box makes for good display on the grocer's shelves.

Padlocks, as an Irish friend puts it, are always getting themselves lost. One of the simplest ways I've seen to keep track of padlocks about the fruit farm is to "lock" them in the packing shed. This is done by hanging a six-inch length of chain over a nail on the wall; then lock the padlocks in the chain links.

CARBOY CART AND TILTER •

Dumping acid out of a full carboy for use in the fruit washer is a pretty ticklish business unless you have the right equipment. Commercial hydrochloric acid used in the washers will burn if it gets on the skin, and it eats away clothes if spilled on them. New type carboy carts and tilters are now available that take the danger out of handling carboys. The cart allows for moving carboys about the packing shed easily, and with the tilter you can pour acid out of the carboy with ease whether the container is full or almost empty. Since the wood casing on all carboys is the same size, all carboys will fit in the cart rack and tilter.

LOOK OUT ANTI-FREEZE-
HERE I COME!



NOW'S THE TIME TO CLEAN OUT YOUR RADIATOR

PUT your truck in shape for summer driving, right now. Remove anti-freeze and clean out all the rust and sediment that are clogging your radiator. You can do it yourself for ten cents—in a few minutes. Just get a can of Sani-Flush.

Pour a little in the radiator (directions are on the can). Run the motor. Drain. Flush. Then refill with clean water. Sani-Flush removes lime and scale from the cooling system. Opens the circulation of water. Insures an efficient flow through the cooling system. Sani-Flush cannot injure aluminum cylinder heads or motor fittings. Most bathrooms have Sani-Flush for cleaning toilets. Sold at grocery, drug, hardware, or five-and-ten-cent stores—25 and 10 cent sizes. The Hygienic Products Co., Canton, Ohio.

Sani-Flush *Safe*
KEEPS RADIATORS CLEAN NOT CAUSTIC


WATER PUMPS WATER

from spring, creek or artesian well having at least 2-ft. fall and flowing 2 gallons per minute or more. A Rife Ram pumps water to house and barn, or for irrigation. Continuous flow. No operating cost. Money's worth or money back. Send for catalog.

RIFE RAM & PUMP WORKS
BOX 17, WAYNESBORO, VA.



PRESS APPLES FOR PROFIT



A HYDRAULIC CIDER PRESS

will put you into business making good money... big demand for apple cider, grape and fruit juices. Turn undergrade apples into money.

A. B. FARQUHAR CO., Limited Box 13, York, Pa.

Standard Garden Tractors 1 & 2 CYLINDERS

Powerful 1 and 2 Cylinder Tractors for Small Farms, Gardeners, Florists, Nurseries, Fruit and Poultry Men.

NEW MODELS
With Ample Power for Field, Hay and Truck Crop Tools.
Also Run Belt Machines, Pumps, Saws, etc. High Wheels-Enclosed Gears.

LOW PRICES
Write for Easy Terms Plan and **Free Catalog**

STANDARD ENGINE CO.
Minneapolis, Minn. Philadelphia, Pa. New York, N. Y.
3211 Como Ave. 3445 Market St. 308 Cedar St.

SUCCESSFUL ORCHARDS

● A "ROUND TABLE" PAGE FOR EVERY GROWER ●

BEES BECOME APARTMENT DWELLERS

FRUIT GROWERS will soon be giving thought to bees. Some may be wondering if their colonies are strong enough for proper pollination. Ralph R. Van Dine of New York, tells of his experience with some bees that were *too* strong.

"Perhaps some of your readers of the 'Round Table' page who keep bees might be interested to know what a swarm of bees can do to make trouble.

"I have a relatively small orchard of 70 trees and for pollination purposes I have five strong colonies of Italian bees.

"Because I happen to know a little about bees, the president of our bee society and several beekeepers asked me to accompany them to the north side of Long Island to rid a garage-apartment of eight strong—extra strong—colonies of bees.

"It so happened that five years ago someone lost a swarm of bees. The queen of the swarm located a garage with chauffeur's apartments above it. Above each window was a small gable which made a nice place for the hive. The bees got into eight of these gables and all through the walls and the chauffeur and his family had to move out.

"It took more than a month to get the bees out of the walls with smoke. All of the gables had to be removed and holes had to be made in the plaster through which to inject the smoke. Several pounds of honey were removed from the inside of the gables, walls, and living rooms. But this was certainly a costly way to get honey!

"Each swarm was made up of thousands of bees. The swarms weighed from seven to nine pounds each. The house was a mess and it is pretty certain that somebody got stung.

"So, if any 'Round Table' readers who have bees, fail to have the queen's wings clipped, they had better watch out when swarming time comes or the bees might go up in the storage loft, the house attic or walls, or in any of the buildings. After the bees are located in any of these places for awhile, with queens laying 1000 to 3000 or more eggs a day, somebody's going to have plenty of bees."

And your "Round Table" editor might add: That same somebody is going to have lots of trouble. Clipping of queen's wings is not so difficult if the person doing it knows his business, and it helps to keep your bees in the orchard. How about some more ideas on bees? Send them in for the "Round Table" page.

GRAFTING SUCCESSFUL FOR MICHIGAN GROWER

MICHIGAN fruit grower, Glen H. Pilkinton has enjoyed exceptional success with bridge grafting and sends in the following:

"I have found many ideas in your magazine and am sending you an item which might be helpful to other growers. Here is a suggestion for bridge grafting.

"We cut the suckers or scions from trees and store them in a cold place until time for use, after freezing weather. The

This page is a place for growers to get together and exchange experiences and ideas. The beginner, as well as the veteran, will find here many practical suggestions for better and more profitable fruit growing. In return for the helps you receive from this page, be ready to pass on, for the benefit of others, any new idea, method or procedure you have developed or run across. Just jot it down as it occurs to you (a postcard will often do) and mail it to the "ROUND TABLE EDITOR," AMERICAN FRUIT GROWER. Don't worry about fancy writing. What the readers of this page want are practical pointers—that are to the point.

trunk of the girdled tree is covered with soil or a wax. When the time comes for grafting, the soil is removed from around the trunk. The ends

of the scion are cut in a similar manner as when cleft grafting. One of the cut ends of the scion is placed on the bark at the edge of the girdle and a "V" is cut with the knife back-slanted slightly towards the center of the "V". This cuts the scion and the bark at the same time. Remove wood from the thin side of the scion, wedge if necessary to make the scion the same thickness as the bark. Then force the scion into the "V" with the thick part out. Measure the scion, allowing enough length to span girdle, and repeat process at upper end. A scion with bow in it, preferably one that grew on the side of the limb, is best. Place tacks so the heads will cover the ends of the scions in the "V" of the bark.

"The grafts are spaced two to three inches apart around the trunk. I use

AMERICAN FRUIT GROWER

brush or black wax. The grafts are then covered with a burlap sack held in place by a tack at the lap and the sack is covered with soil. I have grafted 40 or 50 trees in my orchard by this method and have not lost a tree. I am sending sample of graft to show more clearly how the unions are made."

Accompanying this item is an illustration showing the sample of bridge grafting sent in by Mr. Pilkinton.

COLOR GUARDS AID IN KEEPING RECORD

FROM James P. Kegel, Jr., Wisconsin fruit grower, comes a new idea in keeping a record of orchard trees. A record of the trees is important for every fruit farm and "Round Table" readers will be interested in what Mr. Kegel has to say.

"I use the following method to keep an accurate record of the variety and age of replacement trees: Each year after cutting out the wire rabbit guards from the roll, I stack the squares over a large pan, placing heavy weights on each corner of the pile. Then I pour over the pile a quantity of colored implement paint thinned a bit to make it flow easier. A different color each year or for each variety helps me identify the stock after the guards have been placed."

Your "Round Table" editor will be glad to have more ideas on orchard record keeping from readers.


UNIQUE EXPERIENCE IN TOP-WORKING TREES

PENNSYLVANIA fruit grower, J. Russell Edgerton, writes regarding his experience with top-working fruit trees:

"In looking over some old copies of AMERICAN FRUIT GROWER I noted a comment on the 'Successful Orchards' page to the effect that completely top-working a tree in one operation was apt to be fatal. I know that is the most common understanding, but some observations have led me to wonder whether there may not be a variable in the picture in the form of soil condition or a similar factor.

"Some years ago I ruthlessly dehorned a 15-year-old open center tree at about two feet from the center crotches, set two or more scions in each stub, and let nature do her best. She did so remarkably well that every scion grew and that tree a dozen years later can scarcely be told from its neighbors except that it is better shaped.

"Two years ago we worked over some less vigorous and older trees on the same soil and chose to be more conservative in how much we did at one time. The result was growth so weak that it is debatable whether to continue the project or not. There were a few places cut back much more than others, to a degree similar to the earlier job, and these scions all grew vigorously. A further setting on more wood last spring gave the same result, so now I am planning to cut back to strong-growing scions and see whether they 'spruce up' or 'pass out.'"



A NEW, ECONOMICAL WAY
TO MAKE SULFUR WETTABLE

WETTABLE

CUT YOUR SUMMER FUNGICIDE SPRAY BILL **ONE-THIRD**

Join the thousands of interested growers who are going to cut their scab-control spray bill **ONE-THIRD** this summer by using the NEW Sherwin-Williams Dry Lime Sulfur-325 Mesh Sulfur—Wettable combination instead of expensive wettable sulfurs. Mix this new Sherwin-Williams combination in your own tank and you get a scab-controlling, non-russetting, non-injurious-to-foliage spray that **SPREADS AND STICKS**.

ASSURES FINE FINISH—FINE COLOR

This newly devised combination which is arousing interest and endorsement everywhere it is demonstrated will protect your crop against scab, against

the risk of russetting from the use of Liquid Lime Sulfur, against foliage injury—and assures **FINE FINISH—FINE COLOR**. Study the Micro-Camera photographs below and you will understand why.

CONTROL CHERRY LEAF SPOT WITH S-W BASI-COP

Here's a New, More Economical and Better-Than-Bordeaux method of controlling cherry leaf spot. S-W Basi-Cop protects your cherry crop without injury to fruit or foliage. It's better for 7 reasons than old-style Bluestone.




WRITE FOR FREE FOLDER THAT TELLS WHY!

THE SHERWIN-WILLIAMS Co.

Insecticide Department

101 Prospect Ave.

Cleveland, Ohio



THE MICRO-CAMERA TELLS THE STORY


Left—Cortland apple leaf sprayed with 2 lbs. of Dry Lime Sulfur and 8 lbs. of 325 mesh sulfur to 100 gals. of water. Note heavy, uniform deposit of colloidal, flocculated sulfur.

Center—Same leaf after 1/2 inch rain. Note good coating of sulfur still remaining. Right—Micro-photograph of surface sprayed with 2 lbs. of Dry Lime Sulfur and 8 lbs. of 325

mesh sulfur to 100 gals. of water. The Dry Lime Sulfur in this combination has flocculated the sulfur and has also acted as a "tooth," holding the sulfur particles tenaciously to the surface.

SHERWIN-WILLIAMS

SPRAY AND DUST MATERIALS



A BANE TO PESTS—A BOON TO GROWERS

In a single season Dow "MIKE" Sulfur made a name for itself.

Introduced last year, this better wettable sulfur was received by growers with real enthusiasm.

This year, orchardists in every fruit-growing state look to the control of brown rot and apple scab with greater confidence for they know the results they can expect with "MIKE" Sulfur on the job.

Dow "MIKE" Sulfur is 15 times finer than 325 mesh sulfur with a particle size of approximately 1/5000 of an inch. Such fineness results in a dense, fog-like spray that gives exceptional coverage and protection.

And, Dow chemists have made a big advance in adhesion. Once "MIKE" is on the fruit and foliage it stays put for a longer time—even under the washing action of rain. Does not injure foliage.

"MIKE" provides extremely high toxicity. It is more than 95 per cent active sulfur.

Finally, it is far easier to use. It goes into suspension with amazing ease and speed—even without the aid of an agitator. And it stays in suspension longer.

No wonder fruit growers hailed "MIKE" as a real boon to orchard control.

Put "MIKE" to work in your orchard now. Get a trial order. See how it mixes—how it sprays—how completely it covers—how long it lasts—and what positive protection it provides. Just one experience with "MIKE" and you will never go back to ordinary sulfurs.

INSTANTLY WETTABLE • HIGHER TOXICITY
GREATER ADHESION • DISTINCT ECONOMY
MICROSCOPIC FINENESS

MAKE THE DOW "MIKE" SULFUR TABLE TEST

Write for a free sample of Dow "MIKE" Sulfur. See for yourself how fine it is.

Then, make this test. Pour the sample into a glass of water. See how fast it goes into suspension—how long it stays that way.

Let it settle overnight. Just shake the glass and see how it goes back into suspension. What it does before your eyes it does in actual spraying practice.

The sample is FREE—no obligation—just write for it today.



THE DOW CHEMICAL COMPANY, MIDLAND, MICHIGAN

Branch Sales Offices: 30 Rockefeller Plaza, New York City—Second and Madison Streets, St. Louis—135 South La Salle Street, Chicago